Index of Authors

ABELSON, H. T., SPECTOR, R., GORKA, C. & FOSBURG, M. Kinetics of tetrahydrobiopterin synthesis by rabbit brain dihydrofolate reductase 267–268

ALI, S. Y. *see* BAYLISS, M. T. 149–154

ALLEN, A. K., DESAI, N. N., NEUBERGER, A. & CREETH, J. M. Properties of potato lectin and the nature of its glycoprotein linkages 665–674

ANDERSEN, J. A. *see* BROOKS, R. R. 725–732

ARONSON, N. N. & BARRETT, A. J. The specificity of cathepsin B. Hydrolysis of glucagon at the *C*-terminus by a peptidyldipeptidase mechanism 759–765

ARZABE, F. R. *see* BROCK, J. H. 73–78

ASHBY, P., TOLSON, A. M. & ROBINSON, D. S. The heterogeneity of the lipoprotein lipase of rat epididymal adipose tissue 305–311

ASPART, L., COOKE, R., DELSENY, M. & GUITTON, Y. Effect of the protein-synthesis-initiation inhibitor 2-(4-methyl-2,6-dinitroanilino)-*N*-methylpropionamide on ribonucleic acid synthesis in radish seedlings 607–611

ATASSI, M. Z. & LEE, C.-L. Boundary refinement of the lysozyme antigenic site around the disulphide bond 6–127 (site 1) by 'surface-simulation' synthesis 419–427

ATASSI, M. Z. & LEE, C.-L. The precise and entire antigenic structure of native lysozyme 429–434

AURICCHIO, F., ROTONDI, A., SAMPAOLO, P. & SCHIAVONE, E. Cytosol oestrogen receptor of lactating mammary gland. Effect of heparin on the aggregation of the receptor and interaction of the receptor with heparin–Sepharose 137–141

AVILA, J. L. The influence of the type of sulphate bond and degree of sulphation of glycosaminoglycans on their interaction with lysosomal enzymes 489–491

BAIG, M. A. & SALAHUDDIN, A. Occurrence and characterization of stable intermediate state(s) in the unfolding of ovomucoid by guanidine hydrochloride 89–97

BANERJEE, C. K. *see* SANI, B. P. 711–717

BARBER, M. J. *see* BRAY, R. C. 653–658

BARKER, G. R. & HARDMAN, N. The effects of Acridine Orange on deoxyribonucleic acid in *Escherichia coli* 567–573

BARRETT, A. J. *see* ARONSON, N. N. 759–765

BAYLISS, M. T. & ALI, S. Y. Studies on cathepsin B in human articular cartilage 149–154

BEGUIN, C. G. *see* GOUT, E. 719–723

BELL, M. V. *see* HACKING, A. J. 41–50

BILLINGS, R. E. *see* TUKEY, R. H. 659–663

BLACKWELL, L. F. *see* MACGIBBON, A. K. H. 527–531; 533–538

BLANCKAERT, N., COMPERNOLLE, F., LEROY, P., VAN HOUTTE, R., FEVERY, J. & HEIRWEGH, K. P. M. The fate of bilirubin-IXα glucuronide in cholestasis and during storage *in vitro*. Intramolecular rearrangement to positional isomers of glucuronic acid 203–214

BLANCKAERT, N. *see also* COMPERNOLLE, F. 185–201

BLUARD-DECONINCK, J.-M., WILLIAMS, J., EVANS, R. W., VAN SNICK, J., OSINSKI, P. A. & MASSON, P. L. Iron-binding fragments from the *N*-terminal and *C*-terminal regions of human lactoferrin 321–327

BRAY, R. C., BARBER, M. J. & LOWE, D. J. Electron-paramagnetic-resonance spectroscopy of complexes of xanthine oxidase with xanthine and uric acid 653–658

BRAY, R. C. *see also* VINCENT, S. P. 639–647

BRINDLEY, D. N. *see* STURTON, R. G. 263–266

BROCK, J. H., ARZABE, F. R., RICHARDSON, N. E. & DEVERSON, E. V. Characterization of monoferric fragments obtained by tryptic cleavage of bovine transferrin 73–78

BROCKLEHURST, K. *see* SHIPTON, M. 385–401

BROOKS, R. R. & ANDERSEN, J. A. Substrate, metal and template effects on inhibition of bacteriophage-Qβ ribonucleic acid polymerase by ortho- and pyrophosphate 725–732

BRZOZOWSKI, A. M. *see* GRABOWSKI, M. J. 277–279

BUCKLEY, P. D. *see* MACGIBBON, A. K. H. 527–531; 533–538

BURCHELL, B. & HALLINAN, T. Phospholipid content and activity of pure uridine diphosphate–glucuronyltransferase from rat liver. 821–824

CAMMACK, R., HUCKLESBY, D. P. & HEWITT, E.J. Electron-paramagnetic-resonance studies of the mechanism of leaf nitrite reductase. Signals from the iron–sulphur centre and haem under turnover conditions 519–526

CAMPBELL, A. M. Conformational variation in superhelical deoxyribonucleic acid 281–283

CAPUTTO, R. *see* MAGGIO, B. 559–565

CHANCE, B., SARONIO, C., LEIGH, JR., J. S., INGLEDEW, W. J. & KING, T. E. Low-temperature kinetics of the reaction of oxygen and solubilized cytochrome oxidase 787–798

CHARTERIS, A. *see* JOHN, R. A. 771–779

CHESNE, S. *see* GOUT, E. 719–723

CLAYCOMB, W. C. Biochemical aspects of cardiac muscle differentiation. Determination of deoxyribonucleic acid template availability and 3′-hydroxyl termini in nuclei and chromatin by using exogenous deoxyribonucleic acid polymerases 289–298

COATES, J. H. *see* HARDMAN, M. J. 215–223

COBURN, S. P. & SCHALTENBRAND, W. E. Simplified preparation of a phosphatase inhibitor and further studies of its action 485–488

COLBY, J. & DALTON, H. Resolution of the methane monooxygenase of *Methylococcus capsulatus* (Bath) into three components. Purification and properties of component C, a flavoprotein 461–468

COMPERNOLLE, F., VAN HEES, G. P., BLANCKAERT, N. & HEIRWEGH, K. P. M. Glucuronic acid conjugates of bilirubin-IXα in normal bile compared with post-obstructive bile. Transformation of the 1-*O*-acylglucuronide into 2-, 3- and 4-*O*-acylglucuronides 185–201

COMPERNOLLE, F. *see also* BLANCKAERT, N. 203–214

COOKE, R. *see* ASPART, L. 607–611

CORTIJO, M., JIMENEZ, J. S. & LLOR, J. Criteria to recognize the structure and micropolarity of pyridoxal 5′-phosphate-binding sites in proteins 497–500

COUGHLAN, M. P. *see* NÍ FHAOLÁIN, Í. 83–88

CRABTREE, B. *see* TAYLOR, D. J. 273–275

CREETH, J. M. *see* ALLEN, A. K. 665–674

CRICHTON, R. R., PONCE-ORTIZ, Y., KOCH, M. H. J., PARFAIT, R. & STUHRMANN, H. B. Isolation and characterization of phytoferritin from pea (*Pisum sativum*) and lentil (*Lens esculenta*) 349–356
CUMAR, F. A. *see* MAGGIO, B. 559–565
CUMMINS, P. & PERRY, S. V. Troponin I from human skeletal and cardiac muscles 251–259
CYGLER, M. *see* GRABOWSKI, M. J. 277–279

DABEVA, M. D., DUDOV, K. P., HADJIOLOV, A. A. & STOYKOVA, A. S. Quantitative analysis of rat liver nucleolar and nucleoplasmic ribosomal ribonucleic acids 367–374
DABEVA, M. D. *see also* DUDOV, K. P. 375–383
DAHLMANN, B. & REINAUER, H. Purification and some properties of an alkaline proteinase from rat skeletal muscle 803–810
DALTON, H. *see* COLBY, J. 461–468
DALZIEL, K., MCFERRAN, N., MATTHEWS, B. & REYNOLDS, C. H. Transient kinetics of nicotinamide–adenine dinucleotide phosphate-linked isocitrate dehydrogenase from bovine heart mitochondria 743–750
DALZIEL, K. *see also* REYNOLDS, C. H. 733–742
DELANEY, M. E. & WALKER, D. A. Comparison of the kinetic properties of ribulose biphosphate carboxylase in chloroplast extracts of spinach, sunflower and four other reductive pentose phosphate-pathway species 477–482
DELSENY, M. *see* ASPART, L. 607–611
DEREWENDA, A. E. *see* GRABOWSKI, M. J. 277–279
DEREWENDA, Z. A. *see* GRABOWSKI, M. J. 277–279
DESAI, N. N. *see* ALLEN, A. K. 665–674
DEVERSON, E. V. *see* BROCK, J. H. 73–78
D'HONTD, A. *see* D'HONDT, J. 705–710
D'HONDT, J., MOENS, L., HEIP, J., D'HONDT, A. & KONDO, M. Oxygen-binding characteristics of three extracellular haemoglobins of *Artemia salina* 705–710
DICKENSON, C. J. & DICKINSON, F. M. Inhibition by ethanol, acetaldehyde and trifluoroethanol of reactions catalysed by yeast and horse liver alcohol dehydrogenase 613–627
DICKENSON, C. J. *see also* DICKINSON, F. M. 629–637
DICKINSON, F. M. & DICKENSON, C. J. Estimation of rate and dissociation constants involving ternary complexes in reactions catalysed by yeast alcohol dehydrogenase 629–637
DICKINSON, F. M. *see also* DICKENSON, C. J. 613–627
DICKSON, I. R. & ROUGHLEY, P. J. A comparative study of the proteoglycan of growth cartilage of normal and rachitic chicks 675–682
DIXON, G. H. *see* GEDAMU, L. 589–599
DUDOV, K. P., DABEVA, M. D., HADJIOLOV, A. A. & TODOROV, B. N. Processing and migration of ribonsomal ribonucleic acids in the nucleolus and nucleoplasm of rat liver nuclei 375–383
DUDOV, K. P. *see also* DABEVA, M. D. 367–374
DUGAL, B. Effect of different compounds on 1-aspartamido-β-*N*-acetylglucosamine amidohydrolase from human liver 799–802
DUGGLEBY, R. G. *see* ELLIS, K. J. 513–517

EDGERTON, M. E. *see* MOORE, T. A. 469–476
EISEN, H. J. & GLINSMANN, W. H. Maximizing the purification of the activated glucocorticoid receptor by DNA–cellulose chromatography 177–183
ELLIS, K. J. & DUGGLEBY, R. G. What happens when data are fitted to the wrong equation? 513–517
ENGEL, P. C. & JONES, J. B. A reappraisal of the reaction of butyryl-coenzyme A dehydrogenase with phenylmercuric acetate. Evidence that de-greening involves a reaction of the tightly bound thioester 51–59
EVANS, R. W. *see* BLUARD-DECONINCK, J.-M. 321–327

FARRAR, G. H. & HARRISON, R. Isolation and structural characterization of alkali-labile oligosaccharides from bovine milk-fat-globule membrane 549–557
FERENCZI, M. A., HOMSHER, E., SIMMONS, R. M. & TRENTHAM, D. R. Reaction mechanism of the magnesium ion-dependent adenosine triphosphatase of frog muscle myosin and subfragment 1 165–175
FERENCZI, M. A., HOMSHER, E., TRENTHAM, D. R. & WEEDS, A. G. Preparation and characterization of frog muscle myosin subfragment 1 and actin 155–163
FEVERY, J. *see* BLANCKAERT, N. 203–214
FLODGAARD, H. & TORP-PEDERSEN, C. A calcium ion-dependent adenosine triphosphate pyrophosphohydrolase in plasma membrane from rat liver. Demonstration that the adenosine triphosphate analogues adenosine 5′-[$\beta\gamma$-imido]triphosphate and adenosine 5′-[$\beta\gamma$-methylene]triphosphate are substrates for the enzyme 817–820
FOSBURG, M. *see* ABELSON, H. T. 267–268
FOWLER, L. J. *see* JOHN, R. A. 771–779

GAHMBERG, C. G., VIRTANEN, I. & WARTIOVAARA, J. Cross-linking of erythrocyte membrane proteins by periodate and intramembrane particle distribution 683–686
GALDES, A. & HILL, H. A. O. The proton exchange of the *pro-S* hydrogen atom at C-1 in dihydroxyacetone phosphate and D-fructose 1,6-bisphosphate catalysed by class-I and class-II aldolases 539–542
GEDAMU, L., IATROU, K. & DIXON, G. H. A simple procedure for the isolation and purification of protamine messenger ribonucleic acid from trout testis 589–599
GLINSMANN, W. H. *see* EISEN, H. J. 177–183
GLOVER, G. I. *see* MARIANO, P. S. 115–122
GODDARD, J. P. *see* LOWDON, M. 601–606
GOLDBERG, A. & LEHMAN, W. Troponin-like proteins from muscles of the scallop, *Aequipecten irradians* 413–418
GORKA, C. *see* ABELSON, H. T. 267–268
GOUT, E., CHESNE, S., BEGUIN, C. G. & PELMONT, J. Kinetic studies with the use of proton-magnetic-resonance spectroscopy of the specific α-deuteration of amino acids by *Escherichia coli* aspartate aminotransferase 719–723
GRABOWSKI, M. J., BRZOZOWSKI, A. M., DEREWENDA, Z. S., SKARŻYŃSKI, T., CYGLER, M., STĘPIEŃ, A. & DEREWENDA, A. E. Crystallization of human oxyhaemoglobin from poly(ethylene glycol) solutions 277–279
GREENWOOD, C. *see* MOORE, T. A. 469–476
GUITTON, Y. *see* ASPART, L. 607–611
GUTFREUND, H. *see* HARDMAN, M. J. 215–223

HACKING, A. J., BELL, M. V. & HASSALL, H. The purification and properties of urocanase from *Pseudomonas testosteroni* 41–50
HADJIOLOV, A. A. *see* DABEVA, M. D. 367–374; DUDOV, K. P. 375–383
HALLINAN, T. *see* BURCHELL, B. 821–824
HAMBERG, U. *see* VAHTERA, E. 767–770
HARDING, J. J. *see* PANJWANI, N. A. 687–695; 697–703
HARDINGHAM, T. E. *see* SHEEHAN, J. K. 109–114
HARDMAN, M. J., COATES, J. H. & GUTFREUND, H. Pressure relaxation of the equilibrium of the pig heart lactate dehydrogenase system 215–223
HARDMAN, N. *see* BARKER, G. R. 567–573
HARRIS, W. J. *see* ROSS, C. A. 231–240; 241–249
HARRISON, P. M. *see* TREFFRY, A. 313–320
HARRISON, R. *see* FARRAR, G. H. 549–557
HASLEWOOD, G. A. D., IKAWA, S., TÖKÉS, L. & WONG, D. Bile salts of the green turtle *Chelonia mydas* (L) 409–412
HASSALL, H. *see* HACKING, A. J. 41–50
HAYLOCK, S. J. *see* MACGIBBON, A. K. H. 533–538
HEIP, J. *see* D'HONDT, J. 705–710
HEIRWEGH, K. P. M. *see* BLANCKAERT, N. 203–214; COMPERNOLLE, F. 185–201
HERMANN, R. L. *see* NEUMANN, J. R. 123–135
HEWITT, E. J. *see* CAMMACK, R. 519–526
HILL, H. A. O. *see* GALDES, A. 539–542
HOMSHER, E. *see* FERENCZI, M. A. 155–163; 165–175
HOOD, K. & ROBERTS, G. C. K. Ultraviolet difference-spectroscopic studies of substrate and inhibitor binding to *Lactobacillus casei* dihydrofolate reductase 357–366
HUCKLESBY, D. P. *see* CAMMACK, R. 519–526
HUGHES, D. G. & MADEN, B. E. H. The pseudouridine contents of the ribosomal ribonucleic acids of three vertebrate species. Numerical correspondence between pseudouridine residues and 2′-*O*-methyl groups is not always conserved 781–786
HYNES, M. J. *see* NÍ FHAOLÁIN, Í. 83–88

IATROU, K. *see* GEDAMU, L. 589–599
IKAWA, S. *see* HASLEWOOD, G. A. D. 409–412
INGLEDEW, W. J. *see* CHANCE, B. 787–798
IUPAC–IUB Commission on Biochemical Nomenclature. Nomenclature of multiple forms of enzymes (Recommendations 1976) 37–39
IUPAC–IUB Commission on Biochemical Nomenclature. Nomenclature of phosphorus-containing compounds of biochemical importance (Recommendations 1976) 1–19
IUPAC–IUB Commission on Biochemical Nomenclature. The nomenclature of lipids (Recommendations 1976) 21–35

JACOBSEN, C. Lysine residue 240 of human serum albumin is involved in high-affinity binding of bilirubin 453–459
JIMENEZ, J. S. *see* CORTIJO, M. 497–500
JOHN, R. A., CHARTERIS, A. & FOWLER, L. J. The reaction of amino-oxyacetate with pyridoxal phosphate-dependent enzymes 771–779
JOHNSON, A. R. *see* WARD, P. E. 143–148
JONES, J. B. *see* ENGEL, P. C. 51–59

KEIGHREN, M. A. & PRICE, N. C. A study of the role of the reactive thiol group of rabbit muscle creatine kinase with a chromophoric reporter group 269–272
KERR, M. A. & PORTER, R. R. The purification and properties of the second component of human complement 99–107
KING, T. E., *see* CHANCE, B. 787–798
KOCH, M. H. J. *see* CRICHTON, R. R. 349–356
KONDO, M. *see* D'HONDT, J. 705–710
KUCHEL, P. W. *see* REYNOLDS, C. H. 733–742

LAWLEY, P. D. *see* SWENSON, D. H. 575–587
LEE, C.-L. *see* ATASSI, M. Z. 419–427; 429–434
LEHMAN, W. *see* GOLDBERG, A. 413–418
LEIGH, JR., J. S. *see* CHANCE, B. 787–798
LEROY, P. *see* BLANCKAERT, N. 203–214
LLOR, J. *see* CORTIJO, M. 497–500
LO, H.-S. & REEVES, R. E. Pyruvate-to-ethanol pathway in *Entamoeba histolytica* 225–230
LOWDON, M. & GODDARD, J. P. Chemical modification as a probe of conformational changes in transfer ribonucleic acid on aminoacylation 601–606
LOWE, D. J. Electron paramagnetic resonance in biochemistry. Computer simulation of spectra from frozen aqueous samples 649–651
LOWE, D. J. *see also* BRAY, R. C. 653–658

MACGIBBON, A. K. H., BLACKWELL, L. F. & BUCKLEY, P. D. Steady-state and pre-steady-state kinetic studies on mitochondrial sheep liver aldehyde dehydrogenase. A comparison with the cytoplasmic enzyme 527–531
MACGIBBON, A. K. H., HAYLOCK, S. J., BUCKLEY, P. D. & BLACKWELL, L. F. Kinetic studies on the esterase activity of cytoplasmic sheep liver aldehyde dehydrogenase 533–538
MADEN, B. E. H. *see* HUGHES, D. G. 781–786
MAGGIO, B., CUMAR, F. A. & CAPUTTO, R. Surface behaviour of gangliosides and related glycosphingolipids 559–565
MARIANO, P. S., GLOVER, G. I. & PETERSEN, J. R. Evidence for a pH-dependent irreversible formation of a stable conformation of phenacyl-α-chymotrypsin 115–122
MASSON, P. L. *see* BLUARD-DECONINCK, J.-M. 321–327
MATTHEWS, B. *see* DALZIEL, K. 743–750
MAYER, H. *see* THARANATHAN, R. N. 403–408
MCCRAE, S. I. *see* WOOD, T. M. 61–72
MCFERRAN, N. *see* DALZIEL, K. 743–750
MEIKLE, M. C. *see* SELLERS, A. 493–496
MOENS, L. *see* D'HONDT, J. 705–710
MOORE, T. A., EDGERTON, M. E., PARR, G., GREENWOOD, C. & PERHAM, R. N. Studies of an acid-induced species of purple membrane from *Halobacterium halobium* 469–476
MUIR, H. *see* SHEEHAN, J. K. 109–114

NEUBERGER, A. *see* ALLEN, A. K. 665–674
NEUMANN, J. R., O'MEARA, A. R. & HERRMANN, R. L. Cyclic AMP-dependent histone-specific nucleoplasmic protein kinase from rat liver 123–135
NÍ FHAOLÁIN, Í., HYNES, M. J. & COUGHLAN, M. P. Stopped-flow spectrophotometric studies on the reaction of turkey liver xanthine dehydrogenase with reducing substrates 83–88
NIEDUSZYNSKI, I. A. *see* SHEENAN, J. K. 109–114

OFFER, G. *see* STARR, R. 813–816
O'MEARA, A. R. *see* NEUMANN, J. R. 123–135

OSINSKI, P. A. *see* BLUARD-DECONINCK, J.-M. 321–327
OZA, N. B. & RYAN, J. W. A simple high-yield procedure for isolation of human urinary kallikreins 285–288

PANJWANI, N. A. & HARDING, J. J. Chymotryptic and tryptic peptides of fragment α_1-CB3 from bovine corneal collagen. Pinpointing the sites of hexose attachment 697–703
PANJWANI, N. A. & HARDING, J. J. Isolation and hydroxylysine glycoside content of some cyanogen bromide-cleaved fragments of collagen from bovine corneal stroma 687–695
PARFAIT, R. *see* CRICHTON, R. R. 349–356
PARKER, P. J. & RANDLE, P. J. Partial purification and properties of branched-chain 2-oxo acid dehydrogenase of ox liver 751–757
PARR, G. *see* MOORE, T. A. 469–476
PELMONT, J. *see* GOUT, E. 719–723
PERHAM, R. N. *see* MOORE, T. A. 469–476
PERRY, S. V. *see* CUMMINS, P. 251–259
PETERSEN, J. R. *see* MARIANO, P. S. 115–122
PHELPS, C. F. *see* SHEEHAN, J. K. 109–114
PONCE-ORTIZ, Y. *see* CRICHTON, R. R. 349–356
PORTER, R. R. *see* KERR, M. A. 99–107
PRICE, N. C. *see* KEIGHREN, M. A. 269–272
PROPPING, D., ZANEVELD, L. J. D., TAUBER, P. F. & SCHUMACHER, G. F. B. Purification of plasminogen activators from human seminal plasma 435–444

RANDLE, P. J. *see* PARKER, P. J. 751–757
REEVES, R. E. *see* LO, H.-S. 225–230
REINAUER, H. *see* DAHLMANN, B. 803–810
REYNOLDS, C. H., KUCHEL, P. W. & DALZIEL, K. Equilibrium binding of coenzymes and substrates to nicotinamide–adenine dinucleotide phosphate-linked isocitrate dehydrogenase from bovine heart mitochondria 733–742
REYNOLDS, C. H. *see also* DALZIEL, K. 743–750
REYNOLDS, J. J. *see* SELLERS, A. 493–496
RICHARDSON, N. E. *see* BROCK, J. H. 73–78
ROBERTS, G. C. K. *see* HOOD, K. 357–366
ROBINSON, D. S. *see* ASHBY, P. 305–311
ROSS, C. A. & HARRIS, W. J. DNA polymerases from *Chlamydomonas reinhardii*. Further characterization, action of inhibitors and associated nuclease activities 241–249
ROSS, C. A. & HARRIS, W. J. DNA polymerases from *Chlamydomonas reinhardii*. Purification and properties 231–240
ROTONDI, A. *see* AURICCHIO, F. 137–141
ROUGHLEY, P. J. *see* DICKSON, I. R. 675–682
RYAN, J. W. *see* OZA, N. B. 285–288

SALAHUDDIN, A. *see* BAIG, M. A. 89–97
SAMPAOLO, P. *see* AURICCHIO, F. 137–141
SANDERSON, A. R. *see* TURNER, M. J. 505–508
SANI, B. P., TITUS, B. C. & BANERJEE, C. K. Determination of binding affinities of retinoids to retinoic acid-binding protein and serum albumin 711–717
SARONIO, C. *see* CHANCE, B. 787–798
SCHALTENBRAND, W. E. *see* COBURN, S. P. 485–488
SCHIAVONE, E. *see* AURICCHIO, F. 137–141
SCHIEBEL, W. *see* ZÄNKER, K. S. 445–451
SCHUMACHER, G. F. B. *see* PROPPING, D. 435–444
SELLERS, A., REYNOLDS, J. J. & MEIKLE, M. C. Neutral metallo-proteinases of rabbit bone. Separation in latent forms of distinct enzymes that when activated degrade collagen, gelatin and proteoglycans 493–496
SHEEHAN, J. K., NIEDUSZYNSKI, I. A., PHELPS, C. F., MUIR, H. & HARDINGHAM, T. E. Self-association of proteoglycan subunits from pig laryngeal cartilage 109–114
SHIPTON, M. & BROCKLEHURST, K. Characterization of the papain active centre by using two-protonic-state electrophiles as reactivity probes. Evidence for nucleophilic reactivity in the un-interrupted cysteine-25–histidine-159 interactive system 385–401
SIMMONS, R. M. *see* FERENCZI, M. A. 165–175
SKARŻYŃSKI, T. *see* GRABOWSKI, M. J. 277–279
SMITH, G. H. *see* TAYLOR, D. J. 273–275
SMITH, G. P. Sequence of the full-length immunoglobulin κ-chain of mouse myeloma MPC 11 337–347
SNOSWELL, A. M. & TUBBS, P. K. Deacylation of acetylcoenzyme A and acetylcarnitine by liver preparations 299–303
SPECTOR, R. *see* ABELSON, H. T. 267–268
STARR, R. & OFFER, G. The interaction of C-protein with heavy meromyosin and subfragment-2 813–816
STĘPIEŃ, A. *see* GRABOWSKI, M. J. 277–279
STORRING, P. L. & TIPLADY, R. J. A simple sensitive method, applicable to small polypeptides, for the determination of proteins in polyacrylamide gels after isoelectric focusing 79–82
STOYKOVA, A. S. *see* DABEVA, M. D. 367–374
STUHRMANN, H. B. *see* CRICHTON, R. R. 349–356
STURTON, R. G. & BRINDLEY, D. N. Problems encountered in measuring the activity of phosphatidate phosphohydrolase 263–266
SUGIMOTO, K. *see* TOMODA, A. 329–335
SUHARA, M. *see* TOMODA, A. 329–335
SWENSON, D. H. & LAWLEY, P. D. Alkylation of deoxyribonucleic acid by carcinogens dimethyl sulphate, ethyl methanesulphonate, *N*-ethyl-*N*-nitrosourea and *N*-methyl-*N*-nitrosourea. Relative reactivity of the phosphodiester site thymidylyl(3′-5′)thymidine 575–587

TAKESHITA, M. *see* TOMODA, A. 329–335
TAUBER, P. F. *see* PROPPING, D. 435–444
TAYLOR, D. J., CRABTREE, B. & SMITH, G. H. The intracellular location of pyruvate carboxylase, citrate synthase and 3-hydroxyacyl-CoA dehydrogenase in lactating rat mammary gland 273–275
TEPHLY, T. R. *see* TUKEY, R. H. 659–663
THARANATHAN, R. N., MAYER, H. & WECKESSER, J. Location of *O*-methyl sugars in antigenic (lipo-)polysaccharides of photosynthetic bacteria and cyanobacteria 403–408
TIPLADY, R. J. *see* STORRING, P. L. 79–82
TITUS, B. C. *see* SANI, B. P. 711–717
TODOROV, B. N. *see* DUDOV, K. P. 375–383
TÖKÉS, L. *see* HASLEWOOD, G. A. D. 409–412
TOLSON, A. M. *see* ASHBY, P. 305–311
TOMODA, A., SUGIMOTO, K., SUHARA, M., TAKESHITA, M. & YONEYAMA, Y. Haemichrome formation from haemoglobin subunits by hydrogen peroxide 329–335
TORP-PEDERSEN, C. *see* FLODGAARD, H. 817–820
TREFFRY, A. & HARRISON, P. M. Incorporation and release of inorganic phosphate in horse spleen ferritin 313–320

TRENTHAM, D. R. *see* FERENCZI, M. A. 155–163; 165–175
TUBBS, P. K. *see* SNOSWELL, A. M. 299–303
TUKEY, R. H., BILLINGS, R. E. & TEPHLY, T. R. Separation of oestrone UDP-glucuronyltransferase and *p*-nitrophenol UDP-glucuronyltransferase activities 659–663
TURNER, M. J. & SANDERSON, A. R. The preparation of liposomes bearing human (HLA) transplantation antigens 505–508

VAHTERA, E. & HAMBERG, U. Interaction of urokinase with α_2-macroglobulin investigated by isoelectric focusing. Evidence for non-specific dissociable binding 767–770
VAN HEES, G. P. *see* COMPERNOLLE, F. 185–201
VAN HOUTTE, R. *see* BLANCKAERT, N. 203–214
VAN SNICK, J. *see* BLUARD-DECONINCK, J.M. 321–327
VINCENT, S. P. & BRAY, R. C. Electron-paramagnetic-resonance studies on nitrate reductase from *Escherichia coli* K12 639–647
VIRTANEN, I. *see* GAHMBERG, C. G. 683–686
VLADUTIU, G. D. I-cell disease. A hypothesis for the structure of the carbohydrate recognition site on β-D-*N*-actylhexosaminidase 509–512

WALKER, D. A. *see* DELANEY, M. E. 477–482
WARD, P. E. & JOHNSON, A. R. Renal inactivation of substance P in the rat 143–148
WARTIOVAARA, J. *see* GAHMBERG, C. G. 683–686
WECKESSER, J. *see* THARANATHAN, R. N. 403–408
WEEDS, A. G. *see* FERENCZI, M. A. 155–163
WHITEHEAD, E. P. Co-operativity and the methods of plotting binding and steady-state kinetic data 501–504
WILLIAMS, J. *see* BLUARD-DECONINCK, J.-M. 321–327
WONG, D. *see* HASLEWOOD, G. A. D. 409–412
WOOD, T. M. & MCCRAE, S. I. The cellulase of *Trichoderma kongignii*. Purification and properties of some endoglucanase components with special reference to their action on cellulose when acting alone and in synergism with the cellobiohydrolase 61–72

YONEYAMA, Y. *see* TOMODA, A. 329–335

ZANEVELD, L. J. D. *see* PROPPING, D. 435–444
ZÄNKER, K. S. & SCHIEBEL, W. The heterogeneity of cytoplasmic deoxyribonucleic acid polymerase from *Physarum polycephalum* 445–451

Index of Subjects

Acetaldehyde, inhibition by ethanol, trifluoroethanol and, of reactions catalysed by yeast and horse liver alcohol dehydrogenases (Dickenson, C. J. & Dickinson, F. M.) 613–627

Acetylcarnitine, deacylation of acetyl-coenzyme A and, by rat and sheep liver preparations (Snoswell, A. M. & Tubbs, P. K.) 299–303

Acetyl-coenzyme A, deacylation of acetylcarnitine and, by rat and sheep liver preparations (Snoswell, A. M. & Tubbs, P. K.) 299–303

β-D-*N*-Acetylhexosaminidase, fibroblast, human, hypothesis for the structure of the carbohydrate recognition site on, and its relevance to the aetiology of I-cell disease (Vladutiu, G. D.) 509–512

Acridine Orange, effects of, on deoxyribonucleic acid during growth of *Escherichia coli* (Barker, G. R. & Hardman, N.) 567–573

Actin, skeletal-muscle, frog, preparation and characterization of, and myosin subfragment 1 (Ferenczi, M.A., Homsher, E., Trentham, D. R. & Weeds, A. G.) 155–163

Adenosine 5′-[βγ-imido]triphosphate, demonstration that adenosine 5′-[βγ-methylene]triphosphate and, are substrates for rat liver plasma-membrane calcium ion-dependent adenosine triphosphate pyrophosphatase (Flodgaard, H. & Torp-Pedersen, C.) 817–820

Adenosine 5′-[βγ-methylene]triphosphate, demonstration that adenosine 5′-[βγ-imido]triphosphate and, are substrates for rat liver plasma-membrane calcium ion-dependent adenosine triphosphate pyrophosphatase (Flodgaard, H. & Torp-Pedersen, C.) 817–820

Adenosine triphosphatase, magnesium ion-dependent, myosin and myosin subfragment 1, skeletal-muscle, frog, reaction mechanism of (Ferenczi, M. A., Homsher, E., Simmons, R. M. & Trentham, D. R.) 165–175

Adenosine triphosphate pyrophosphatase, calcium ion-dependent, plasma-membrane, liver, rat, demonstration that adenosine 5′-[βγ-imido]triphosphate and adenosine 5′-[βγ-methylene]triphosphate are substrates for (Flodgaard, H. & Torp-Pedersen, C.) 817–820

Adipose tissue, epididymal, rat, heterogeneity of the lipoprotein lipase from (Ashby, P., Tolson, A. M. & Robinson, D. S.) 305–311

Adrenocorticotropin, *see* Corticotropin

Aequipecten irradians, *see* Scallop

Albumin, serum, human, involvement of lysine residue 240 of, in the high-affinity binding of bilirubin (Jacobsen, C.) 453–459

Albumin, serum, ox, criteria for recognition of the structure and micropolarity of pyridoxal 5′-phosphate-binding sites in, and other proteins (Cortijo, M., Jimenez, J. S. & Llor, J.) 497–500

Albumin, serum, ox, determination of binding affinities of retinoids to chick-embryo skin retinoic acid-binding protein and to (Sani, B. P., Titus, B. C. & Banerjèe, C. K.) 711–717

Alcohol dehydrogenases, yeast and horse liver, inhibition by ethanol, acetaldehyde and trifluoroethanol of reactions catalysed by (Dickenson, C. J. & Dickinson, F. M.) 613–627

Alcohol dehydrogenase, yeast, estimation of rate and dissociation constants involving ternary complexes in reactions catalysed by (Dickinson, F. M. & Dickenson, C. J.) 629–637

Aldehyde dehydrogenase, cytoplasmic, liver, sheep, kinetic properties of the esterase activity of (MacGibbon A. K. H., Haylock, S. J., Buckley, P. D. & Blackwell, L. F.) 533–538

Aldehyde dehydrogenases, mitochondrial and cytoplasmic liver, sheep, comparison of the steady-state and pre-steady-state kinetic properties of (MacGibbon, A. K. H., Blackwell, L. F. & Buckley, P. D.) 527–531

Aldolase, *see* Fructose bisphosphate aldolase

Alkaline proteinase, *see* proteinase, alkaline

Amino acid sequence, determination of, of the full-length κ light chain of mouse myeloma MPC 11 immunoglobulin G (Smith, G. P.) 337–347

Aminoacylation, use of chemical modification as a probe of conformational changes during, of *Escherichia coli* transfer ribonucleic acid (Lowdon, M. & Goddard, J. P.) 601–606

4-Aminobutyrate aminotransferase, brain, rabbit, reaction of amino-oxyacetate with, and with other pyridoxal 5′-phosphate-dependent enzymes (John, R. A., Charteris, A. & Fowler, L. J.) 771–779

Amino-oxyacetate, reaction of, with pyridoxal 5′-phosphate-dependent enzymes (John, R. A., Charteris, A. & Fowler, L. J.) 771–779

Anabaena variabilis, location of *O*-methyl sugars in antigenic lipopolysaccharides and polysaccharides of various photosynthetic bacteria and cyanobacteria including (Tharanathan, R. N., Mayer, H. & Weckesser, J.) 403–408

Anacystis nidulans, location of *O*-methyl sugars in antigenic lipopolysaccharides and polysaccharides of various photosynthetic bacteria and cyanobacteria including (Tharanathan, R. N., Mayer, H. & Weckesser, J.) 403–408

Angiotensinase, role of kininase and, in the inactivation of substance P in rat kidney (Ward, P. E. & Johnson, A. R.) 143–148

Antigenic site 1, lysozyme, egg-white, chicken, boundary refinement by surface-simulation synthesis of, around disulphide bridge 6–127 (Atassi, M. Z. & Lee, C.-L.) 419–427

Antigenic sites 1, 2 and 3, lysozyme, egg-white, chicken, precise structure of (Atassi, M. Z. & Lee, C.-L.) 429–434

Antigens, transplantation, human (HLA), incorporation of, into liposomes (Turner, M. J. & Sanderson, A. R.) 505–508

Apoferritin, spleen, horse, incorporation of iron and inorganic phosphate into (Treffry, A. & Harrison, P. M.) 313–320

Artemia salina, *see* Brine shrimp

Aspartate aminotransferase, *Escherichia coli* K12, kinetic studies with the use of proton-magnetic-resonance spectroscopy of the specific α-deuteration of aspartate and phenylalanine catalysed by (Gout, E., Chesne, S., Beguin, C. G. & Pelmont, J.) 719–723

Aspartate aminotransferase, heart, pig, reaction of amino-oxyacetate with, and with other pyridoxal 5′-phosphate-dependent enzymes (John, R. A., Charteris, A. & Fowler, L. J.) 771–779
Aspartate, kinetic studies with the use of proton-magnetic-resonance spectroscopy of the specific α-deuteration of phenylalanine and, catalysed by *Escherichia coli* K12 aspartate aminotransferase (Gout, E., Chesne, S., Beguin, C. G. & Pelmont, J.) 719–723
Aspartylglucosylaminase, liver, human, effects of various compounds on the activity of (Dugal, B.) 1978, 799–802

Bacteriophage PM2, conformational variation in super-helical deoxyribonucleic acid from (Campbell, A. M.) 281–283
Bacteriophage Qβ, effects of substrate, metal ions and template on the inhibition by inorganic phosphate and inorganic pyrophosphate of the activity of ribonucleic acid polymerase from (Brooks, R. R. & Andersen, J. A.) 725–732
Bacteriorhodopsin, purple-membrane, *Halobacterium halobium*, properties of an acid-induced spectral species of (Moore, T. A., Edgerton, M. E., Parr, G., Greenwood C. & Perham, R. N.) 469–476
Bile, human and rat, intramolecular rearrangement of the 1-*O*-acyl isomer of bilirubin-IXα glucuronide to the 2-, 3- and 4-*O*-acyl isomers in, in cholestasis and during storage *in vitro* (Blanckaert, N., Compernolle, F., Leroy, P., Van Houtte, R., Fevery, J. & Heirwegh, K. P. M.) 203–214
Bile, human and rat, post-obstructive, transformation of the 1-*O*-acyl isomer of bilirubin-IXα glucuronide into the 2-, 3- and 4-*O*-acyl isomers in (Compernolle, F., Van Hees, G. P., Blanckaert, N. & Heirwegh, K. P. M.) 185–201
Bile salts, green-turtle, characterization of (Haslewood, G. A. D., Ikawa, S., Tökés, L. & Wong, D.) 409–412
Bilirubin, involvement of lysine residue 240 of human serum albumin in the high-affinity binding of (Jacobsen, C.) 453–459
Bilirubin-IXα glucuronide, intramolecular rearrangement of the 1-*O*-acyl isomer of, to the 2-, 3- and 4-*O*-acyl isomers in human and rat bile in cholestasis and during storage *in vitro* (Blanckaert, N., Compernolle, F., Leroy, P., Van Houtte, R., Fevery, J. & Heirwegh, K. P. M.) 203–214
Bilirubin-IXα glucuronide, transformation of the 1-*O*-acyl isomer of, into the 2-, 3- and 4-*O*-acyl isomers in post-obstructive human and rat bile (Compernolle, F., Van Hees, G. P., Blanckaert, N. & Heirwegh, K. P. M.) 185–201
Bone, rabbit, separation from, of latent forms of collagenase and of two neutral metallo-proteinases that when activated degrade gelatin and cartilage proteoglycans (Sellers, A., Reynolds, J. J. Meikle, M. C.) 493–496
Brain, rabbit, kinetics of the biosynthesis of tetrahydrobiopterin by tetrahydrofolate dehydrogenase from (Abelson, H. T., Spector, R., Gorka, C. & Fosburg, M.) 267–268
Brain, rabbit, reaction of amino-oxyacetate with 4-aminobutyrate aminotransferase from, and with other pyridoxal 5′-phosphate-dependent enzymes (John, R. A., Charteris, A. & Fowler, L. J.) 771–779
Brine shrimp (*Artemia salina*), oxygen-binding characteristics of three extracellular haemoglobins from (D'hondt, J., Moens, L., Heip, J., D'hondt, A. & Kondo, M.) 705–710
Butyryl-coenzyme A dehydrogenase, *Megasphaera elsdenii*, reappraisal of the reaction of phenylmercuric acetate with (Engel, P. C. & Jones, J. B.) 51–59

Cartilage, articular, human, activity of cathepsin B in (Bayliss, M. T. & Ali, S. Y.) 149–154
Cartilage, growth, chick, effect of vitamin D deficiency on the proteoglycans of (Dickson, I. R. & Roughley, P. J.) 675–682
Cartilage, laryngeal, pig, self-association of subunits of proteoglycans from (Sheehan, J. K., Nieduszynski, I. A., Phelps, C. F., Muir, H. & Hardingham, T. E.) 109–114
Cartilage, separation from rabbit bone of latent forms of collagenase and of two neutral metallo-proteinases that when activated degrade gelatin and proteoglycans from (Sellers, A., Reynolds, J. J. & Meikle, M. C.) 493–496
Cathepsin B, activity of, in human articular cartilage (Bayliss, M. T. & Ali, S. Y.) 149–154
Cathepsin B, lysosomal, liver, human, hydrolysis of glucagon at the *C*-terminus by a peptidyldipeptidase mechanism catalysed by, and by the enzyme from other mammalian sources (Aronson, N. N. & Barrett, A. J.) 759–765
Cellobiohydrolase, cellulase-complex, *Trichoderma koningii* I.M.I. 73022, purification and properties of four *Trichoderma koningii* I.M.I. 73022 cellulase-complex endoglucanases with special reference to their action on cellulose when acting alone and in synergism with (Wood, T. M. & McCrae, S. I.) 61–72
Cells, cultured, evidence for lack of exact numerical correspondence between the contents of pseudouridylate residues and 2′-*O*-methyl groups of ribosomal ribonucleic acids from, of three vertebrate species (Hughes, D. G. & Maden, B. E. H.) 781–786
Cellulase complex, *Trichoderma koningii* I.M.I. 73022, purification and properties of four endoglucanase components of, with special reference to their action on cellulose when acting alone and in synergism with the cellobiohydrolase component (Wood, T. M. & McCrae, S. I.) 61–72
Cellulose, purification and properties of four endoglucanase components of *Trichoderma koningii* I.M.I. 73022 cellulase complex with special reference to their action on, when acting alone and in synergism with cellobiohydrolase component (Wood, T. M. & McCrae, S. I.) 61–72
Chelonia mydas, *see* Green turtle
Chick, effect of vitamin D deficiency on the proteoglycans of growth cartilage of (Dickson, I. R. & Roughley, P. J.) 675–682
Chick-embryo skin, determination of binding affinities of retinoids to ox serum albumin and to retinoic acid-binding protein from (Sani, B. P., Titus, B. C. & Banerjee, C. K.) 711–717

Chicken egg, occurrence and characterization of stable intermediate state(s) in the unfolding by guanidinium chloride of ovomucoid from (Baig, M. A. & Salahuddin, A.) 89–97
Chicken egg-white, boundary refinement by surface-simulation synthesis of the antigenic site (site 1) around disulphide bridge 6–127 of lysozyme from (Atassi, M. Z. & Lee, C.-L.) 419–427
Chicken egg-white, precise structure of all three antigenic sites (sites 1, 2 and 3) of lysozyme from (Atassi, M. Z. & Lee, C.-L.) 429–434
Chlamydomonas reinhardii, characterization of, action of inhibitors on and association of deoxyribonuclease activity with deoxyribonucleic acid polymerases from (Ross, C. A. & Harris, W. J.) 241–249
Chlamydomonas reinhardii, purification and properties of deoxyribonucleic acid polymerases from (Ross, C. A. & Harris, W. J.) 231–240
Chloroplasts, comparison of the kinetic properties of ribulose bisphosphate carboxylase in extracts of, from six higher-plant species utilizing the reductive pentose phosphate pathway (Delaney, M. E. & Walker, D. A.) 477–482
Cholestasis, intramolecular rearrangement of the 1-*O*-acyl isomer of bilirubin IXα glucuronide to the 2-, 3- and 4-*O*-acyl isomers in human and rat bile in, and during storage *in vitro* (Blanckaert, N., Compernolle, F., Leroy, P., Van Houtte, R., Fevery, J. & Heirwegh, K. P. M.) 203–214
Chromatin, heart-muscle, rat, determination of the template activity and 3′-hydroxy-terminal nucleotide residues of deoxyribonucleic acids in, and nuclei by using exogenous deoxyribonucleic acid polymerases (Claycomb, W. C.) 289–298
Chromatium vinosum, location of *O*-methyl sugars in antigenic lipopolysaccharides and polysaccharides of various photosynthetic bacteria and cyanobacteria including (Tharanathan, R. N., Mayer, H. & Weckesser, J.) 403–408
α-Chymotrypsin, phenacyl-, evidence for a pH-dependent irreversible formation of a stable conformation of (Mariano, P. S., Glover, G. I. & Petersen, J. R.) 115–122
Citrate synthase, intracellular location of pyruvate decarboxylase, 3-hydroxyacyl-coenzyme A dehydrogenase and, in lactating rat mammary gland (Taylor, D. J., Crabtree, B. & Smith, G. H.) 273–275
Clearing-factor lipase, *see* Lipoprotein lipase
Collagen, corneal-stroma, ox, identification of the sites containing hydroxylysine glycoside residues in chymotryptic and tryptic peptides of fragment α1-CB3 of (Panjwani, N. A. & Harding, J. J.) 697–703
Collagen, corneal-stroma, ox, isolation and content of hydroxylysine glycoside residues of some cyanogen bromide-cleaved fragments of (Panjwani, N. A. & Harding, J. J.) 687–695
Collagenase, bone, rabbit, separation of latent forms of, and of two neutral metallo-proteinases that when activated degrade gelatin and cartilage proteoglycans (Sellers, A., Reynolds, J. J. & Meikle, M. C.) 493–496
Complement, human, purification and properties of component 2 of (Kerr, M. A. & Porter, R. R.) 99–107
Component 2, complement, human, purification and properties of (Kerr, M. A. & Porter, R. R.) 99–107
Computers, use of, for simulation of electron-paramagnetic-resonance spectra from frozen aqueous samples (Lowe, D. J.) 649–651
Corneal stroma, ox, identification of the sites containing hydroxylysine glycoside residues in chymotryptic and tryptic peptides of fragment α1-CB3 of collagen from Panjwani, N. A. & Harding, J. J.) 697–703
Corneal stroma, ox, isolation and content of hydroxylysine glycoside residues of some cyanogen bromide-cleaved fragments of collagen from (Panjwani, N. A. & Harding, J. J.) 687–695
Corticotropin, pig, simple sensitive method for the determination of, and other proteins and polypeptides in polyacrylamide gels after isoelectric focusing (Storring, P. L. & Tiplady, R. J.) 79–82
Creatine kinase, skeletal-muscle, rabbit, use of a chromophoric reporter group to study the role of the reactive thiol group of (Keighren, M. A. & Price, N. C.) 269–272
Cucurbita pepo, *see* Vegetable marrow
Cysteine residue, active-site, use of two-protonic-state electrophiles as reactivity probes for, of papain (Shipton, M. & Brocklehurst, K.) 385–401
Cytochrome *c* oxidase, heart, ox, solubilized, low-temperature kinetics of the reaction of oxygen with (Chance, B., Saronio, C., Leigh, Jr., J. S., Ingledew, W. J. & King, T. E.) 787–798
Cytosol, liver, rat, maximization of the purification of glucocorticoid–glucocorticoid-receptor protein complex from, by deoxyribonucleic acid–cellulose chromatography (Eisen, H. J. & Glinsmann, W. H.) 177–183
Cytosol, liver, sheep, comparison of the steady-state and pre-steady-state kinetic properties of aldehyde dehydrogenases from, and mitochondria (MacGibbon, A. K. H., Blackwell, L. F. & Buckley, P. D.) 527–531
Cytosol, liver, sheep, kinetic properties of the esterase activity of aldehyde dehydrogenase from (MacGibbon, A. K. H., Haylock, S. J., Buckley, P. D. & Blackwell, L. F.) 533–538
Cytosol, mammary-gland, mouse, lactating, effect of heparin on the aggregation of oestrogen-receptor protein from, and interaction of the receptor protein with heparin–Sepharose (Auricchio, F., Rotondi, A., Sampaolo, P. & Schiavone, E.) 137–141

Data, use of residual plots as a guide in the choice of more appropriate equations when, have been fitted to the wrong equation (Ellis, K. J. & Duggleby, R. G.) 513–517
Deoxyribonuclease, association of activity of, with *Chlamydomonas reinhardii* deoxyribonucleic acid polymerases (Ross, C. A. & Harris, W. J.) 241–249
Deoxyribonucleic acid, bacteriophage PM2, superhelical, conformational variation in (Campbell, A. M.) 281–283
Deoxyribonucleic acid–cellulose, maximization of the purification of rat liver cytosol glucocorticoid–glucocorticoid-receptor protein complex by chromatography on (Eisen, H. J. & Glinsmann, W. H.) 177–183
Deoxyribonucleic acid, effects of Acridine Orange on, during growth of *Escherichia coli* (Barker, G. R. & Hardman, N.) 567–573

Deoxyribonucleic acid polymerase, cytoplasmic, *Physarum polycephalum*, heterogeneity of (Zänker, K. S. & Schiebel, W.) 445–451
Deoxyribonucleic acid polymerases, *Chlamydomonas reinhardii*, characterization of, action of inhibitors on and association of deoxyribonuclease activity with (Ross, C. A. & Harris, W. J.) 241–249
Deoxyribonucleic acid polymerases, *Chlamydomonas reinhardii*, purification and properties of (Ross, C. A. & Harris, W. J.) 231–240
Deoxyribonucleic acid polymerases, exogenous, determination of the template activity and 3′-hydroxy-terminal nucleotide residues of rat heart-muscle nuclear and chromatin deoxyribonucleic acids by using (Claycomb, W. C.) 289–298
Deoxyribonucleic acid, sperm, salmon, relative reactivity of the phosphodiester site thymidylyl(3′-5′)thymidine of, towards alkylation by dimethyl sulphate, ethyl methanesulphonate, *N*-ethyl-*N*-nitrosourea and *N*-methyl-*N*-nitrosourea (Swenson, D. H. & Lawley, P. D.) 575–587
Deoxyribonucleic acids, nuclear and chromatin, heart-muscle, rat, determination of the template activity and 3′-hydroxy-terminal nucleotide residues of, by using exogenous deoxyribonucleic acid polymerases (Claycomb, W. C.) 289–298
Dihydrofolate reductase, *see* Tetrahydrofolate dehydrogenase
Dihydroxyacetone phosphate, exchange of protons involving C-1 *pro-S* hydrogen atoms of fructose 1,6-bisphosphate and, catalysed by class-I and class-II fructose bisphosphate aldolases (Galdes, A. & Hill, H. A. O.) 539–542
3,4-Dihydroxyphenylalanine decarboxylase, kidney, pig, reaction of amino-oxyacetate with, and with other pyridoxal 5′-phosphate-dependent enzymes (John, R. A. Charteris, A. & Fowler, L. J.) 771–779
Dimethyl sulphate, relative reactivity of the phosphodiester site thymidylyl(3′-5′)thymidine of salmon sperm deoxyribonucleic acid towards alkylation by, and other carcinogens (Swenson, D. H. & Lawley, P. D.) 575–587
2,2′-Dipyridyl disulphide, use of n-propyl 2-pyridyl disulphide and, as reactivity probes for the active-site cysteine residue of papain (Shipton, M. & Brocklehurst, K.) 385–401
Disulphide bridge 6–127, boundary refinement by surface-simulation synthesis of the antigenic site (site 1) around, of chicken egg-white lysozyme (Atassi, M. Z. & Lee, C.-L.) 419–427
Dopa, *see* 3,4-Dihydroxyphenylalanine

Egg, chicken, occurrence and characterization of stable intermediate state(s) in the unfolding by guanidinium chloride of ovomucoid from (Baig, M. A. & Salahuddin, A.) 89–97
Egg-white, chicken, boundary refinement by surface-simulation synthesis of the antigenic site (site 1) around disulphide bridge 6–127 of lysozyme from (Atassi, M. Z. & Lee, C.-L.) 419–427
Egg-white, chicken, precise structure of all three antigenic sites (sites 1, 2 and 3) of lysozyme from (Atassi, M. Z. & Lee, C.-L.) 429–434
Electron-paramagnetic-resonance spectra, computer simulation of, from frozen aqueous samples (Lowe, D. J.) 649–651
Endoglucanases, cellulase-complex, *Trichoderma koningii* I.M.I. 73022, four, purification and properties of, with special reference to their action on cellulose when acting alone and in synergism with the cellobiohydrolase component (Wood, T. M. & McCrae, S. I.) 61–72
Entamoeba histolytica, pathway for the metabolism of pyruvate to ethanol in (Lo, H.-S. & Reeves, R. E.) 225–230
Enzymes, co-operativity and the methods of plotting data for the binding of ligands and for steady-state kinetics of reactions catalysed by (Whitehead, E. P.) 501–504
Enzymes, nomenclature of multiple forms of (IUPAC-IUB Commission on Biochemical Nomenclature) 37–39
Epididymis, rat, heterogeneity of the adipose-tissue lipoprotein lipase from (Ashby, P., Tolson, A. M. & Robinson, D. S.) 305–311
Erythrocytes, human, cross-linking by periodate of membrane proteins of, and distribution of intramembrane particles (Gahmberg, C. G., Virtanen, I. & Wartiovaara, J.) 683–686
Escherichia coli, effects of Acridine Orange on deoxyribonucleic acid during growth of (Barker, G. R. & Hardman, N.) 567–573
Escherichia coli K12, electron-paramagnetic-resonance studies on nitrate reductase from (Vincent, S. P. & Bray, R. C.) 639–647
Escherichia coli K12, kinetic studies with the use of proton-magnetic-resonance spectroscopy of the specific α-deuteration of aspartate and phenylalanine catalysed by aspartate aminotransferase from (Gout, E., Chesne, S., Beguin, C. G. & Pelmont, J.) 719–723
Escherichia coli, use of chemical modification as a probe of conformational changes during aminoacylation of transfer ribonucleic acid from (Lowdon, M. & Goddard, J. P.) 601–606
Esterase, kinetic properties of the activity of, of sheep liver cytoplasmic aldehyde dehydrogenase (MacGibbon, A. K. H., Haylock, S. J., Buckley, P. D. & Blackwell, L. F.) 533–538
Estradiol, *see* Oestradiol
Estrogens, *see* Oestrogens
Estrone, *see* Oestrone
Ethanol, inhibition by acetaldehyde, trifluoroethanol and, of reactions catalysed by yeast and horse liver alcohol dehydrogenases (Dickenson, C. J. & Dickinson, F. M.) 613–627
Ethanol, pathway for the metabolism of pyruvate to, in *Entamoeba histolytica* (Lo, H.-S. & Reeves, R. E.) 225–230
Ethyl methanesulphonate, relative reactivity of the phosphodiester site thymidylyl(3′-5′)thymidine of salmon sperm deoxyribonucleic acid towards alkylation by, and other carcinogens (Swenson, D. H. & Lawley, P. D.) 575–587
N-Ethyl-*N*-nitrosourea, relative reactivity of the phosphodiester site thymidylyl(3′-5′)thymidine of salmon sperm deoxyribonucleic acid towards alkylation by, and other carcinogens (Swenson, D. H. & Lawley, P. D.) 575–587
Eye, ox, identification of the sites containing hydroxylysine glycoside residues in chymotryptic and tryptic peptides of fragment α1-CB3 of collagen from corneal stroma of (Panjwani, N. A. & Harding, J. J.) 697–703

Eye, ox, isolation and content of hydroxylysine glycoside residues of some cyanogen bromide-cleaved fragments of collagen from corneal stroma of (Panjwani, N. A. & Harding, J. J.) 687–695

Fatty acids, nomenclature of, and other lipids (IUPAC–IUB Commission on Biochemical Nomenclature) 21–35
Ferritin, spleen, horse, incorporation and release of iron and inorganic phosphate in (Treffry, A. & Harrison, P. M.) 313–320
Ferritins, phyto, lentil-seed and pea-seed, isolation and characterization of (Crichton, R. R., Ponce-Ortiz, Y., Koch, M. H. J., Parfait, R. & Stuhrmann, H. B.) 349–356
Fibroblasts, human, hypothesis for the structure of the carbohydrate recognition site on β-D-*N*-acetylhexosaminidase of, and its relevance to the aetiology of I-cell disease (Vladutiu, G. D.) 509–512
Flavin–adenine dinucleotide, identification of, as the prosthetic group of component C of the *Methylococcus capsulatus* methane mono-oxygenase system (Colby, J. & Dalton, H.) 461–468
Folate, use of ultraviolet difference spectroscopy for study of the binding of, and inhibitors to *Lactobacillus casei* tetrahydrofolate dehydrogenase (Hood, K. & Roberts, G. C. K.) 357–366
Frog (*Rana temporaria* and *Rana pipiens*), preparation and characterization of skeletal-muscle myosin subfragment 1 and actin from (Ferenczi, M. A., Homsher, E., Trentham, D. R. & Weeds, A. G.) 155–163
Frog (*Rana temporaria* and *Rana pipiens*), reaction mechanism of skeletal-muscle myosin and myosin subfragment 1 magnesium ion-dependent adenosine triphosphatase from (Ferenczi, M. A., Homsher, E., Simmons, R. M. & Trentham, D. R.) 165–175
Fructose 1,6-bisphosphate, exchange of protons involving C-1 *pro-S* hydrogen atoms of dihydroxyacetone phosphate and, catalysed by class-I and class-II fructose bisphosphate aldolases (Galdes, A. & Hill, H. A. O.) 539–542
Fructose bisphosphate aldolases, class-I and class-II, exchange of protons involving C-1 *pro-S* hydrogen atoms of dihydroxyacetone phosphate and fructose 1,6-bisphosphate catalysed by (Galdes, A. & Hill, H. A. O.) 539–542

Gangliosides, surface behaviour of monolayers of, and of related glycosphingolipids (Maggio, B., Cumar, F. A. & Caputto, R.) 559–565
Gelatin, separation from rabbit bone of latent forms of collagenase and of two neutral metallo-proteinases that when activated degrade cartilage proteoglycans and (Sellers, A., Reynolds, J. J. & Meikle, M. C.) 493–496
Glucagon, hydrolysis of, at the *C*-terminus by a peptidyldipeptidase mechanism catalysed by lysosomal cathepsin B from human liver and other mammalian sources (Aronson, N. N. & Barrett, A. J.) 759–765
Glucanases, endo-, cellulase-complex, *Trichoderma koningii* I.M.I. 73022, four, purification and properties of, with special reference to their action on cellulose when acting alone and in synergism with the cellobiohydrolase component (Wood, T. M. & McCrae, S. I.) 61–72
Glucocorticoid–glucocorticoid-receptor protein complex, cytosol, liver, rat, maximization of the purification of, by deoxyribonucleic acid–cellulose chromatography (Eisen, H. J. & Glinsmann, W. H.) 177–183
Glucocorticoid-receptor protein, cytosol, liver, rat, maximization of the purification of the complex of glucocorticoid and, by deoxyribonucleic acid–cellulose chromatography (Eisen, H. J. & Glinsmann, W. H.) 177–183
Glucuronide conjugate, bilirubin IXα, intramolecular rearrangement of the 1-*O*-acyl isomer of, to the 2-, 3- and 4-*O*-acyl isomers in human and rat bile in cholestasis and during storage *in vitro* (Blanckaert, N., Compernolle F., Leroy, P., Van Houtte, R., Fevery, J. & Heirwegh, K. P. M.) 203–214
Glucuronide conjugate, bilirubin IXα, transformation of the 1-*O*-acyl isomer of, into the 2-, 3- and 4-*O*-acyl isomers in post-obstructive human and rat bile (Compernolle, F., Van Hees, G. P., Blanckaert, N. & Heirwegh, K. P. M.) 185–201
Glycolipids, nomenclature of, and other lipids (IUPAC–IUB Commission on Biochemical Nomenclature) 21–35
Glycoprotein linkages, lectin, potato-tuber, nature of (Allen, A. K., Desai, N. N., Neuberger, A. & Creeth, J. M.) 665–674
Glycoproteins, membrane, milk-fat-globule, cow, isolation and structural characterization of alkali-labile oligosaccharides obtained from (Farrar, G. H. & Harrison, R.) 549–557
Glycosaminoglycans, influence of the type of sulphate bond and degree of sulphation of, on their interaction with human leucocyte lysosomal enzymes (Avila, J. L.) 489–491
Glycoside residues, content of, attached to hydroxylysine residues of some cyanogen bromide-cleaved fragments of ox corneal-stroma collagen (Panjwani, N. A. & Harding, J. J.) 687–695
Glycoside residues, identification of the sites containing attached to hydroxylysine residues in chymotryptic and tryptic peptides of fragments α1-CB3 of ox corneal-stroma collagen (Panjwani, N. A. & Harding, J.J.) 697–703
Glycosphingolipids, surface behaviour of gangliosides and (Maggio, B., Cumar, F. A. & Caputto, R.) 559–565
Green turtle (*Chelonia mydas*), characterization of the bile salts of (Haslewood, G. A. D., Ikawa, S., Tökés, L. & Wong, D.) 409–412
Growth, effects of Acridine Orange on deoxyribonucleic acid during, of *Escherichia coli* (Barker, G. R. & Hardman, N.) 567–573
Guanidinium chloride, occurrence and characterization of stable intermediate state(s) in the unfolding by, of chicken-egg ovomucoid (Baig, M. A. & Salahuddin, A.) 89–97

Haem group, electron-paramagnetic-resonance signals from the iron–sulphur centre and, during turnover of vegetable-marrow leaf nitrate reductase (Cammack, R., Hucklesby, D. P. & Hewitt, E. J.) 519–526
Haemichrome, transformation by hydrogen peroxide of subunits of human haemoglobin into (Tomoda, A., Sugimoto, K., Suhara, M., Takeshita, M. & Yoneyama, Y.) 329–335

Haemoglobin, human, transformation by hydrogen peroxide of subunits of, into haemichrome (Tomoda, A., Sugimoto, K., Suhara, M., Takeshita, M. & Yoneyama, Y.) 329–335

Haemoglobin, oxy-, human, crystallization of, from poly(ethylene glycol) solutions (Grabowski, M. J., Brzozowski, A. M., Derewenda, Z. S., Skarżyński, T., Cygler, M., Stępień, A. & Derewenda, A. E.) 277–279

Haemoglobins, extracellular, brine-shrimp, three, oxygen-binding characteristics of (D'hondt, J., Moens, L., Heip, J., D'hondt, A. & Kondo, M.) 705–710

Halobacterium halobium, properties of an acid-induced spectral species of purple membrane from (Moore, T. A. Edgerton, M. E., Parr, G., Greenwood, C. & Perham, R. N.) 469–476

Heart muscle, human, properties of troponin I from, and skeletal muscle and from the muscles of other mammalian species (Cummins, P. & Perry, S. V.) 251–259

Heart muscle, rat, determination of the template activity and 3′-hydroxy-terminal nucleotide residues of deoxyribonucleic acids in nuclei and chromatin from, by using exogenous deoxyribonucleic acid polymerases (Claycomb, W. C.) 289–298

Heart, ox, equilibrium binding of coenzymes and substrates to mitochondrial nicotinamide–adenine dinucleotide phosphate-linked isocitrate dehydrogenase from (Reynolds, C. H., Kuchel, P. W. & Dalziel, K.) 733–742

Heart, ox, low-temperature kinetics of the reaction of oxygen with solubilized cytochrome *c* oxidase from (Chance, B. Saronio, C., Leigh, Jr., J. S., Ingledew, W. J. & King, T. E.) 787–798

Heart, ox, transient kinetics of the reaction catalysed by mitochondrial nicotinamide–adenine dinucleotide phosphate-linked isocitrate dehydrogenase from (Dalziel, K., McFerran, N., Matthews, B. & Reynolds, C. H.) 743–750

Heart, pig, pressure relaxation of the equilibrium of the reaction catalysed by lactate dehydrogenase from (Hardman, M. J., Coates, J. H. & Gutfreund, H.) 215–223

Heart, pig, reaction of amino-oxyacetate with aspartate aminotransferase from, and with other pyridoxal 5′-phosphate-dependent enzymes (John, R. A., Charteris, A. & Fowler, L. J.) 771–779

HeLa cells, evidence for the lack of exact numerical correspondence between the contents of pseudouridylate residues and 2′-*O*-methyl groups of ribosomal ribonucleic acids from, and other cells from vertebrate species (Hughes, D. G. & Maden, B. E. H.) 781–786

Helianthus annuus, *see* Sunflower

Hen, *see* Chicken

Heparin, effect of, on the aggregation of lactating mouse mammary-gland cytosol oestrogen-receptor protein and interaction of the receptor protein with heparin–Sepharose (Auricchio, F., Rotondi, A., Sampaolo, P. & Schiavone, E.) 137–141

Heparin–Sepharose, effect of heparin on the aggregation of lactating mouse mammary-gland cytosol oestrogen-receptor protein and interaction of the receptor protein with (Auricchio, F., Rotondi, A., Sampaolo, P. & Schiavone, E.) 137–141

Hexose, content of, attached to hydroxylysine residues of some cyanogen bromide-cleaved fragments of ox corneal-stroma collagen (Panjwani, N. A. & Harding, J. J.) 687–695

Hexose, identification of the sites of attachment of, to hydroxylysine residues in chymotryptic and tryptic peptides of fragment α1-CB3 of ox corneal-stroma collagen (Panjwani, N. A. & Harding, J. J.) 697–703

Histones, partial purification and properties of rat liver nucleoplasmic adenosine 3′:5′-cyclic monophosphate-dependent protein kinase specific for the phosphorylation of (Neumann, J. R., O'Meara, A. R. & Herrmann, R. L.) 123–135

Hydrogen atoms, *pro-S*, C-1, exchange of protons involving of dihydroxyacetone phosphate and fructose 1,6-bisphosphate catalysed by class-I and class-II fructose bisphosphate aldolases (Galdes, A. & Hill, H. A. O.) 539–542

Hydrogen ions, exchange of, involving C-1 *pro-S* hydrogen atoms of dihydroxyacetone phosphate and fructose 1,6-bisphosphate catalysed by class-I and class-II fructose bisphosphate aldolases (Galdes, A. & Hill, H. A. O.) 539–542

Hydrogen peroxide, transformation by, of subunits of human haemoglobin into haemichrome (Tomoda, A., Sugimoto, K., Suhara, M., Takeshita, M. & Yoneyama, Y.) 329–335

3-Hydroxyacyl-coenzyme A dehydrogenase, intracellular location of pyruvate decarboxylase, citrate synthase and, in lactating rat mammary gland (Taylor, D. J., Crabtree, B. & Smith, G. H.) 273–275

Hydroxylysine glycoside residues, content of, of some cyanogen bromide-cleaved fragments of ox corneal-stroma collagen (Panjwani, N. A. & Harding, J. J.) 687–695

Hydroxylysine glycoside residues, identification of the sites containing, in chymotryptic and tryptic peptides of fragment α1-CB3 of ox corneal-stroma collagen (Panjwani, N. A. & Harding, J. J.) 697–703

I-cell disease, hypothesis for the structure of the carbohydrate recognition site on human fibroblast β-D-*N*-acetylhexosaminidase and its relevance to the aetiology of (Vladutiu, G. D.) 509–512

Immunoglobulin G, myeloma MPC 11, mouse, determination of the amino acid sequence of the full-length κ light chain of (Smith, G. P.) 337–347

Inorganic phosphate, *see* Phosphate, inorganic

Inorganic pyrophosphate, *see* Pyrophosphate, inorganic

Intramembrane particles, erythrocyte, human, cross-linking by periodate of membrane proteins and distribution of (Gahmberg, C. G., Virtanen, I. & Wartiovaara, J.) 683–686

Iron, characterization of monoferric fragments obtained by tryptic cleavage of ox transferrin and their inability to donate, to rabbit reticulocytes (Brock, J. H., Arzabe, F. R., Richardson, N. E. & Deverson, E. V.) 73–78

Iron, incorporation and release of inorganic phosphate and, in horse spleen ferritin (Treffry, A. & Harrison, P. M.) 313–320

Iron, properties of fragments from *N*-terminal and *C*-terminal regions of human milk lactoferrin binding (Bluard-Deconinck, J.-M., Williams, J., Evans, R. W., Van Snick, J., Osinski, P. A. & Masson, P. L.) 321–327

Iron–sulphur centre, electron-paramagnetic-resonance signals from the haem group and, during turnover of vegetable-marrow leaf nitrite reductase (Cammack, R., Hucklesby, D. P. & Hewitt, E. J.) 519–526
Isocitrate dehydrogenase, nicotinamide–adenine dinucleotide phosphate-linked, mitochondrial, heart, ox, equilibrium binding of coenzymes and substrates to (Reynolds, C. H., Kuchel, P. W. & Dalziel, K.) 733–742
Isocitrate dehydrogenase, nicotinamide–adenine dinucleotide phosphate-linked, mitochondrial, heart, ox, transient kinetics of the reaction catalysed by (Dalziel, K., McFerran, N., Matthews, B. & Reynolds, C. H.) 743–750

Kallikreins, urinary, human, simple high-yield procedure for isolation of (Oza, N. B. & Ryan, J. W.) 285–288
Kidney cells, *Xenopus laevis*, cultured, evidence for lack of exact numerical correspondence between the contents of pseudouridylate residues and 2′-*O*-methyl groups of ribosomal ribonucleic acids from, and other cells from vertebrate species (Hughes, D. G. & Maden, B. E. H.) 781–786
Kidney, pig, reaction of amino-oxyacetate with 3,4-dihydroxyphenylalanine decarboxylase from, and with other pyridoxal 5′-phosphate-dependent enzymes (John, R. A., Charteris, A. & Fowler, L. J.) 771–779
Kidney, rat, inactivation of substance P in (Ward, P. E. & Johnson, A. R.) 143–148
Kininase, role of angiotensinase and, in the inactivation of substance P in rat kidney (Ward, P. E. & Johnson, A. R.) 143–148
Kininogenins, urinary, human, simple high-yield procedure for isolation of (Oza, N. B. & Ryan, J. W.) 285–288

L cells, mouse, evidence for the lack of exact numerical correspondence between the contents of pseudouridylate residues and 2′-*O*-methyl groups of ribosomal ribonucleic acids from, and other cells from vertebrate species (Hughes, D. G. & Maden, B. E. H.) 781–786
Lactate dehydrogenase, heart, pig, pressure relaxation of the equilibrium of the reaction catalysed by (Hardman, M. J., Coates, J. H. & Gutfreund, H.) 215–223
Lactation, effect of heparin on the aggregation of mouse mammary-gland cytosol oestrogen-receptor protein isolated during, and interaction of the receptor protein with heparin–Sepharose (Auricchio, F., Rotondi, A., Sampaolo, P. & Schiavone, E.) 137–141
Lactation, intracellular location of pyruvate decarboxylase, citrate synthase and 3-hydroxyacyl-coenzyme A dehydrogenase in rat mammary gland during (Taylor, D. J., Crabtree, B. & Smith, G. H.) 273–275
Lactobacillus casei, use of ultraviolet difference spectroscopy for study of the binding of substrate and inhibitors to tetrahydrofolate dehydrogenase from (Hood, K. & Roberts, G. C. K.) 357–366
Lactoferrin, milk, human, properties of iron-binding fragments from *N*-terminal and *C*-terminal regions of (Bluard-Deconinck, J.-M., Williams, J., Evans, R. W., Van Snick, J., Osinski, P. O. & Masson, P. L.) 321–327
Larynx, pig, self-association of subunits of proteoglycans from cartilage of (Sheehan, J. K., Nieduszynski, I. A., Phelps, C. F., Muir, H. & Hardingham, T. E.) 109–114
Leaves, comparison of the kinetic properties of ribulose bisphosphate carboxylase in extracts of chloroplasts from, of six higher-plant species utilizing the reductive pentose phosphate pathway (Delaney, M. E. & Walker, D. A.) 477–482
Leaves, vegetable-marrow, electron-paramagnetic-resonance signals from the iron–sulphur centre and haem group during turnover of nitrite reductase from (Cammack, R., Hucklesby, D. P. & Hewitt, E. J.) 519–526
Lectin, potato-tuber, properties of, and the nature of its glycoprotein linkages (Allen, A. K., Desai, N. N., Neuberger, A. & Creeth, J. M.) 665–674
Lens esculenta, *see* Lentil
Lentil (*Lens esculenta*) seeds, isolation and characterization of phytoferritins from pea (*Pisum sativum*) seeds and (Crichton, R. R., Ponce-Ortiz, Y., Koch, M. H. J., Parfait, R. & Stuhrmann, H. B.) 349–356
Leucocytes, human, influence of the type of sulphate bond and degree of sulphation of glycosaminoglycans on their interaction with lysosomal enzymes from (Avila, J. L.) 489–491
Ligands, co-operativity and the methods of plotting data for the binding of, and for steady-state kinetics of reactions catalysed by enzymes (Whitehead, E. P.) 501–504
Light chain, κ, full-length, immunoglobulin G, myeloma MPC 11, mouse, determination of the amino acid sequence of (Smith, G. P.) 337–347
Lipids, nomenclature of (IUPAC–IUB Commission on Biochemical Nomenclature) 21–35
Lipopolysaccharides, location of *O*-methyl sugars in antigenic polysaccharides and, of various photosynthetic bacteria and cyanobacteria (Tharanathan, R. N., Mayer, H. & Weckesser, J.) 403–408
Lipoprotein lipase, adipose-tissue, epididymal, rat, heterogeneity of (Ashby, P., Tolson, A. M. & Robinson, D. S.) 305–311
Liposomes, incorporation of human (HLA) transplantation antigens into (Turner, M. J. & Sanderson, A. R.) 505–508
Liver, horse, inhibition by ethanol, acetaldehyde and trifluoroethanol of reactions catalysed by alcohol dehydrogenases from yeast and (Dickenson, C. J. & Dickinson, F. M.) 613–627
Liver, human, effects of various compounds on the activity of aspartylglucosylaminase from (Dugal, B) 799–802
Liver, human, hydrolysis of glucagon at the *C*-terminus by a peptidyldipeptidase mechanism catalysed by lysosomal cathepsin B from, and other mammalian sources (Aronson, N. N. & Barrett, A. J.) 759–765
Liver, ox, partial purification and properties of mitochondrial branched-chain 2-oxo acid dehydrogenase complex from (Parker, P. J. & Randle, P. J.) 751–757
Liver, rabbit, separation of the activities of microsomal uridine diphosphate glucuronosyltransferases from, specific oestrone and *p*-nitrophenol (Tukey, R. H., Billings, R. E. & Tephly, T. R.) 659–663

Liver, rat and sheep, deacylation of acetyl-coenzyme A and acetylcarnitine by preparations of (Snoswell, A. M. & Tubbs, P. K.) 299–303
Liver, rat, content of phospholipids and activity of microsomal uridine diphosphate glucuronyltransferase purified from (Burchell, B. & Hallinan, T.) 821–824
Liver, rat, demonstration that adenosine 5′-[βγ-imido]-triphosphate and adenosine 5′-[βγ-methylene]triphosphate are substrates for plasma-membrane calcium ion-dependent adenosine triphosphate pyrophosphatase from (Flodgaard, H. & Torp-Pedersen, C.) 817–820
Liver, rat, maximization of the purification of cytosol glucocorticoid–glucocorticoid-receptor protein complex from, by deoxyribonucleic acid–cellulose chromatography (Eisen, H. J. & Glinsmann, W. H.) 177–183
Liver, rat, partial purification and properties of histone-specific nucleoplasmic adenosine 3′:5′-cyclic monophosphate-dependent protein kinase from (Neumann, J. R., O'Meara, A. R. & Herrmann, R. L.) 123–135
Liver, rat, problems encountered in measurement of the activity of phosphatidate phosphatase in microsomal fraction from (Sturton, R. G. & Brindley, D. N.) 263–266
Liver, rat, processing and migration of ribosomal ribonucleic acids in the nucleoli and nucleoplasm of nuclei of (Dudov, K. P., Dabeva, M. D., Hadjiolov, A. A. & Todorov, B. N.) 375–383
Liver, rat, quantitative analysis of nucleolar and nucleoplasmic ribosomal ribonucleic acids from (Dabeva, M. D., Dudov, K. P., Hadjiolov, A. A. & Stoykova, A. S.) 367–374
Liver, sheep, comparison of the steady-state and pre-steady state-kinetic properties of mitochondrial and cytoplasmic aldehyde dehydrogenases from (MacGibbon, A. K. H., Blackwell, L. F. & Buckley, P. D.) 527–531
Liver, sheep, kinetic properties of the esterase activity of cytoplasmic aldehyde dehydrogenase from (MacGibbon, A. K. H., Haylock, S. J., Buckley, P. D. & Blackwell, L. F.) 533–538
Liver, turkey, stopped-flow spectrophotometric studies on the reaction of xanthine dehydrogenase from, with reducing substrates (Ní Fhaoláin, Í., Hynes, M. J. & Coughlan, M. P.) 83–88
Lysine residue 240, involvement of, of human serum albumin in the high-affinity binding of bilirubin (Jacobsen, C.) 453–459
Lysosomes, leucocyte, human, influence of the type of sulphate bond and degree of sulphation of glycosaminoglycans on their interaction with enzymes from (Avila, J. L.) 489–491
Lysosomes, liver, human, hydrolysis of glucagon at the *C*-terminus by a peptidyldipeptidase mechanism catalysed by cathepsin B from, and other mammalian sources (Aronson, N. N. & Barrett, A. J.) 759–765
Lysozyme, egg-white, chicken, boundary refinement by surface-simulation synthesis of the antigenic site (site 1) around disulphide bridge 6–127 of (Atassi, M. Z. & Lee, C.-L.) 419–427
Lysozyme, egg-white, chicken, precise structure of all three antigenic sites (sites 1, 2 and 3) of (Atassi, M. Z. & Lee, C.-L.) 429–434
α_2-Macroglobulin, human, evidence for non-specific dissociable binding during investigation by isoelectric focusing of the interaction of urokinase with (Vahtera, E. & Hamberg, U.) 767–770
Mammary gland, mouse, lactating, effect of heparin on the aggregation of cytosol oestrogen-receptor protein from, and interaction of the receptor protein with heparin–Sepharose (Auricchio, F., Rotondi, A., Sampaolo, P. & Schiavone, E.) 137–141
Mammary gland, rat, lactating, intracellular location of pyruvate carboxylase, citrate synthase and 3-hydroxyacyl-coenzyme A dehydrogenase in (Taylor, D. J., Crabtree, B. & Smith, G. H.) 273–275
Megasphaera elsdenii, reappraisal of the reaction of phenylmercuric acetate with butyryl-coenzyme A dehydrogenase from (Engel, P. C. & Jones, J. B.) 51–59
Membrane, erythrocyte, human, cross-linking by periodate of proteins of, and distribution of intramembrane particles (Gahmberg, C. G., Virtanen, I. & Wartiovaara, J.) 683–686
Membrane, milk-fat-globule, cow, isolation and structural characterization of alkali-labile oligosaccharides obtained from glycoproteins of (Farrar, G. H. & Harrison, R.) 549–557
Membrane, plasma, liver, rat, demonstration that adenosine 5′-[βγ-imido]triphosphate and adenosine 5′-[βγ-methylene]triphosphate are substrates for calcium ion-dependent adenosine triphosphate pyrophosphatase from (Flodgaard, H. & Torp-Pedersen, C.) 817–820
Membrane, purple, *Halobacterium halobium*, properties of an acid-induced spectral species of (Moore, T. A., Edgerton, M. E., Parr, G., Greenwood, C. & Perham, R. N.) 469–476
Messenger ribonucleic acid, *see* Ribonucleic acid, messenger
Methane mono-oxygenase system, *Methylococcus capsulatus*, purification and properties of component C of (Colby, J. & Dalton, H.) 461–468
2-(4-Methyl-2,6-dinitroanilino)-*N*-methylpropionamide, effect of, on the biosynthesis of ribosomal ribonucleic acid in radish seedlings (Aspart, L., Cooke, R., Delseny, M. & Guitton, Y.) 607–611
2′-*O*-Methyl groups, evidence for lack of exact numerical correspondence between the contents of pseudouridylate residues and, of ribosomal ribonucleic acids from cells of three vertebrate species (Hughes, D. G. & Maden, B. E. H.) 781–786
N-Methyl-*N*-nitrosourea, relative reactivity of the phosphodiester site thymidylyl(3′-5′)thymidine of salmon sperm deoxyribonucleic acid towards alkylation by, and other carcinogens (Swenson, D. H. & Lawley, P. D.) 575–587
O-Methyl sugars, location of, in antigenic lipopolysaccharides and polysaccharides of various photosynthetic bacteria and cyanobacteria (Tharanathan, R. N., Mayer, H. & Weckesser, J.) 403–408
Methylococcus capsulatus, purification and properties of component C of the methane mono-oxygenase system from (Colby, J. & Dalton, H.) 461–468
Microsomal fraction, liver, rabbit, separation of the activities of uridine diphosphate glucuronosyltransferases from, specific for oestrone and *p*-nitrophenol (Tukey, R. H., Billings, R. E. & Tephly, T. R.) 659–663

Microsomal fraction, liver, rat, content of phospholipids and activity of uridine diphosphate glucuronyltransferase purified from (Burchell, B. & Hallinan, T.) 821–824

Microsomal fraction, liver, rat, problems encountered in measurement of the activity of phosphatidate phosphatase in (Sturton, R. G. & Brindley, D. N.) 263–266

Milk, electron-paramagnetic-resonance spectroscopy of complexes of xanthine oxidase from, with xanthine and uric acid (Bray, R. C., Barber, M. J. & Lowe, D. J.) 653–658

Milk-fat globules, cow, isolation and structural characterization of alkali-labile oligosaccharides obtained from glycoproteins of the membrane of (Farrar, G. H. & Harrison, R. (549–557)

Milk, human, properties of iron-binding fragments from *N*-terminal and *C*-terminal regions of lactoferrin from (Bluard-Deconinck, J.-M., Williams, J., Evans, R. W., Van Snick, J., Osinski, P. A. & Masson, P. L.) 321–327

Mitochondria, heart, ox, equilibrium binding of coenzymes and substrates to nicotinamide–adenine dinucleotide phosphate-linked isocitrate dehydrogenase from (Reynolds, C. H., Kuchel, P. W. & Dalziel, K.) 733–742

Mitochondria, heart, ox, transient kinetics of the reaction catalysed by nicotinamide-adenine dinucleotide phosphate-linked isocitrate dehydrogenase from (Dalziel, K., McFerran, N., Matthews, B. & Reynolds, C. H.) 743–750

Mitochondria, liver, ox, partial purification and properties of branched-chain 2-oxo acid dehydrogenase complex from (Parker, P. J. & Randle, P. J.) 751–757

Mitochondria, liver, sheep, comparison of the steady-state and pre-steady-state kinetic properties of aldehyde dehydrogenases from, and cytoplasm (MacGibbon, A. K. H., Blackwell, L. F. & Buckley, P. D.) 527–531

Mitochondria, liver, sheep, deacylation of acetyl-coenzyme A by (Snoswell, A. M. & Tubbs, P. K.) 299–303

Mitochondria, mammary-gland, rat, lactating, intracellular location of pyruvate decarboxylase, citrate synthase and 3-hydroxyacylcoenzyme A dehydrogenase in (Taylor, D. J., Crabtree, B. & Smith, G. H.) 273–275

Molybdenum, electron-paramagnetic-resonance spectroscopy of, of milk xanthine oxidase in complexes with xanthine and uric acid (Bray, R. C., Barber, M. J. & Lowe, D. J.) 653–658

Molybdenum, electron-paramagnetic-resonance studies on, of *Escherichia coli* nitrate reductase (Vincent, S. P. & Bray, R. C.) 639–647

Mucolipidosis II, *see* I-cell disease

Muscle, adductor, scallop, preparation and properties of troponin-like proteins from (Goldberg, A. & Lehman, W.) 413–418

Muscle, heart, human, properties of troponin I from, and skeletal muscle and from the muscles of other mammalian species (Cummins, P. & Perry, S. V.) 251–259

Muscle, heart, rat, determination of the template activity and 3′-hydroxy-terminal nucleotide residues of deoxyribonucleic acids in nuclei and chromatin from, by using exogenous deoxyribonucleic acid polymerases (Claycomb, W. C.) 289–298

Muscle, skeletal, frog, preparation and characterization of myosin subfragment 1 and actin from (Ferenczi, M. A., Homsher, E., Trentham, D. R. & Weeds, A. G.) 155–163

Muscle, skeletal, frog, reaction mechanism of myosin and myosin subfragment 1 magnesium ion-dependent adenosine triphosphatase from (Ferenczi, M. A., Homsher, E., Simmons, R. M. & Trentham, D. R.) 165–175

Muscle, skeletal, human, properties of troponin I from, and heart muscle and from the muscles of other mammalian species (Cummins, P. & Perry, S. V.) 251–259

Muscle, skeletal, interaction of C-protein with heavy meromyosin and subfragment 2 of myosin from (Starr, R. & Offer, G.) 813–816

Muscle, skeletal, rabbit, use of a chromophoric reporter group to study the role of the reactive thiol group of creatine kinase from (Keighren, M. A. & Price, N. C.) 269–272

Muscle, skeletal, rat, purification and properties of an alkaline proteinase associated with myofibrillar proteins from (Dahlmann, B. & Reinauer, H.) 803–810

Myeloma MPC 11, mouse, determination of the amino acid sequence of the full-length κ light chain of immunoglobulin G from (Smith, G. P.) 337–347

Myofibrils, skeletal-muscle, rat, purification and properties of an alkaline proteinase associated with proteins of (Dahlmann, B. & Reinauer, H.) 803–810

Myosin, skeletal-muscle, frog, preparation and characterization of subfragment 1 of, and actin (Ferenczi, M. A., Homsher, E., Trentham, D. R. & Weeds, A. G.) 155–163

Myosin, skeletal-muscle, frog, reaction mechanism of magnesium ion-dependent adenosine triphosphatase activity of, and myosin subfragment 1 (Ferenczi, M. A., Homsher, E., Simmons, R. M. & Trentham, D. R.) 165–175

Myosin, skeletal-muscle, interaction of C-protein with heavy meromyosin and subfragment 2 of (Starr, R. & Offer, G.) 813–816

Nicotinamide–adenine dinucleotide, reduced, stopped-flow spectrophotometric studies on the reaction of turkey liver xanthine dehydrogenase with (Ní Fhaoláin, I., Hynes, M. J. & Coughlan, M. P.) 83–88

Nitrate reductase, *Escherichia cloi* K12, electron-paramagnetic-resonance studies on (Vincent, S. P. & Bray, R. C.) 639–647

Nitrite reductase, vegetable-marrow-leaf, electron-paramagnetic-resonance signals from the iron–sulphur centre and haem group during turnover of (Cammack, R., Hucklesby, D. P. & Hewitt, E. J.) 519–526

p-Nitrophenol, separation of the activities of rabbit liver microsomal uridine diphosphate glucuronosyltransferases specific for oestrone and (Tukey, R. H., Billings, R. E. & Tephly, T. R.) 659–663

Nuclei, heart-muscle, rat, determination of the template activity and 3′-hydroxy-terminal nucleotide residues of deoxyribonucleic acids in, and chromatin by using exogenous deoxyribonucleic acid polymerases (Claycomb, W. C.) 289–298

Nuclei, liver, rat, processing and migration of ribosomal ribonucleic acids in the nucleoli and nucleoplasm of (Dudov, K. P., Dabeva, M. D., Hadjiolov, A. A. & Todorov, B. N.) 375–383
Nuclei, liver, rat, quantitative analysis of nucleolar and nucleoplasmic ribosomal ribonucleic acids from (Dabeva, M. D., Dudov, K. P., Hadjiolov, A. A. & Stoykova, A. S.) 367–374
Nucleoli, liver, rat, processing and migration of ribosomal ribonucleic acids in, and nucleoplasm (Dubov, K. P., Dabeva, M. D., Hadjiolov, A. A. & Todorov, B. N.) 375–383
Nucleoli, liver, rat, quantitative analysis of ribosomal ribonucleic acids from, and nucleoplasm (Dabeva, M. D., Dudov, K. P., Hadjiolov, A. A. & Stoykova, A. S.) 367–374
Nucleoplasm, liver, rat, partial purification and properties of histone-specific adenosine 3′:5′-cyclic monophosphate-dependent protein kinase from (Neumann, J. R., O'Meara, A. R. & Herrmann, R. L.) 123–135
Nucleoplasm, liver, rat, processing and migration of ribosomal ribonucleic acids in, and nucleoli (Dudov, K. P., Dabeva, M. D., Hadjiolov, A. A. & Todorov, B. N.) 375–383
Nucleoplasm, liver, rat, quantitative analysis of ribosomal ribonucleic acids from, and nucleoli (Dabeva, M. D., Dudov, K. P., Hadjiolov, A. A. & Stoykova, A. S.) 367–374
Nucleoside triphosphates, nomenclature of analogues of, and other phosphorus-containing compounds of biochemical importance (IUPAC–IUB Commission on Biochemical Nomenclature) 1–19
Nucleotide residues, 3′-hydroxy-terminal, determination of the template activity and, of rat heart-muscle nuclear and chromatin deoxyribonucleic acids by using exogenous deoxyribonucleic acid polymerases (Claycomb, W. C.) 289–298

Oestradiol-17β-receptor protein, cytosol, mammary-gland, mouse, lactating, effect of heparin on the aggregation of, and interaction of the receptor protein with heparin–Sepharose (Auricchio, F., Rotondi, A., Sampaolo, P. & Schiavone, E.) 137–141
Oestrogen-receptor protein, cytosol, mammary-gland, mouse, lactating, effect of heparin on the aggregation of, and interaction of the receptor protein with heparin–Sepharose (Auricchio, F., Rotondi, A., Sampaolo, P. & Schiavone, E.) 137–141
Oestrone, separation of the activities of rabbit liver microsomal uridine diphosphate glucuronosyltransferases specific for *p*-nitrophenol and (Tukey, R. H., Billings, R. E. & Tephly, T. R.) 659–663
Oligosaccharides, alkali-labile, isolation and structural characterization of, obtained from cow milk-fat-globule membrane glycoproteins (Farrar, G. H. & Harrison, R.) 549–557
Osteoarthrosis, activity of cathepsin B in human articular cartilage from normal subjects and from patients with (Bayliss, M. T. & Ali, S. Y.) 149–154
Ovomucoid, chicken-egg, occurrence and characterization of stable intermediate state(s) in the unfolding by guanidinium chloride of (Baig, M. A. & Salahuddin, A.) 89–97
2-Oxo acid (branched-chain) dehydrogenase, mitochondrial, liver, ox, partial purification and properties of (Parker, P. J. & Randle, P. J.) 751–757
Oxygen, characteristics of the binding of, by three brine-shrimp extracellular haemoglobins (D'hondt, J., Moens, L., Heip, J., D'hondt, A. & Kondo, M.) 705–710
Oxygen, low-temperature kinetics of the reaction of, with solubilized ox heart cytochrome *c* oxidase (Chance, B., Saronio, C., Leigh, Jr., J. S., Ingledew, W. J. & King, T. E.) 787–798
Oxyhaemoglobin, human, crystallization of, from poly(ethylene glycol) solutions (Grabowski, M. J., Brzozowski, A. M., Derewenda, Z. S., Skarżyński, T., Cygler, M., Stępień, A. & Derewenda, A. E.) 277–279

Papain, use of two-protonic-state electrophiles as reactivity probes for the active-site cysteine residue of (Shipton, M. & Brocklehurst, K.) 385–401
Pea (*Pisum sativum*) seeds, isolation and characterization of phytoferritins from lentil (*Lens esculenta*) seeds and (Crichton, R. R., Ponce-Ortiz, Y., Koch, M. H. J., Parfait, R. & Stuhrmann, H. B.) 349–356
Pentose phosphate pathway, reductive comparison of the kinetic properties of ribulose bisphosphate carboxylase in extracts of chloroplasts from six higher-plant species utilizing (Delaney, M. E. & Walker, D. A.) 477–482
Peptostreptococcus elsdenii, *see Megasphaera elsdenii*
Periodate, cross-linking by, of human erythrocyte membrane proteins and distribution of intramembrane particles (Gahmberg, C. G., Virtanen, I. & Wartiovaara, J.) 683–686
Peroxide, hydrogen, transformation by, of subunits of human haemoglobin into haemichrome (Tomoda, A., Sugimoto, K., Suhara, M., Takeshita, M. & Yoneyama, Y.) 329–335
Phage, *see* Bacteriophage
Phenacyl-α-chymotrypsin, evidence for a pH-dependent irreversible formation of a stable conformation of (Mariano, P. S., Glover, G. I. & Petersen, J. R.) 115–122
Phenylalanine, kinetic studies with the use of proton-magnetic-resonance spectroscopy of the specific α-deuteration of aspartate and, catalysed by *Escherichia coli* K12 aspartate aminotransferase (Gout, E., Chesne, S., Beguin, C. G. & Pelmont, J.) 719–723
Phenylmercuric acetate, reappraisal of the reaction of, with *Megasphaera elsdenii* butyryl-coenzyme A dehydrogenase (Engel, P. C. & Jones, J. B.) 51–59
Phosphatases, simplified synthesis of 1-pyrrolidinecarbothioic acid (2-pyridylmethylene)hydrazide and its action as an inhibitor of (Coburn, S. P. & Schaltenbrand, W. E.) 485–488
Phosphate, inorganic, effects of substrate, metal ions and template on the inhibition by inorganic pyrophosphate and, of the activity of bacteriophage Qβ-ribonucleic acid polymerase (Brooks, R. R. & Andersen, J. A.) 725–732
Phosphate, inorganic, incorporation and release of iron and, in horse spleen ferritin (Treffry, A. & Harrison, P. M.) 313–320
Phosphatidate phosphatase, problems encountered in measurement of the activity of, in rat liver microsomal fraction (Sturton, R. G. & Brindley, D. N.) 263–266

Phospholipids, content of, and activity of purified rat liver microsomal uridine diphosphate glucuronyltransferase (Burchell, B. & Hallinan, T.) 821–824
Phospholipids, nomenclature of, and other lipids (IUPAC-IUB Commission on Biochemical Nomenclature) 21–35
Phosphorus, nomenclature of compounds of biochemical importance containing (IUPAC–IUB Commission on Biochemical Nomenclature) 1–19
Physarum polycephalum, heterogeneity of cytoplasmic deoxyribonucleic acid polymerase from (Zänker, K. S. & Schiebel, W.) 445–451
Pisum sativum, *see* Pea
Phytoferritins, lentil-seed and pea-seed, isolation and characterization of (Crichton, R. R., Ponce-Ortiz, Y., Koch, M. H. J., Parfait, R. & Stuhrmann, H. B.) 349–356
Plasma membrane, *see* Membrane, plasma
Plasminogen activators, seminal-plasma, human, two, purification and properties of (Propping, D., Zaneveld, L. J. D., Tauber, P. F. & Schumacher, G. F. B.) 435–444
Polypeptides, simple sensitive method for the determination of proteins and, in polyacrylamide gels after isoelectric focusing (Storring, P. L. & Tiplady, R. J.) 79–82
Polysaccharides, location of *O*-methyl sugars in antigenic lipopolysaccharides and, of various photosynthetic bacteria and cyanobacteria (Tharanathan, R. N., Mayer, H. & Weckesser, J.) 403–408
Potato (*Solanum tuberosum*) tubers, properties of the lectin from, and the nature of its glycoprotein linkages (Allen, A. K., Desai, N. N., Neuberger, A. & Creeth, J. M.) 665–674
Pressure, effect of, on the equilibrium of the reaction catalysed by pig heart lactate dehydrogenase (Hardman, M. J., Coates, J. H. & Gutfreund, H.) 215–223
n-Propyl 2-pyridyl disulphide, use of 2,2′-dipyridyl disulphide and, as reactivity probes for the active-site cysteine residue of papain (Shipton, M. & Brocklehurst, K.) 385–401
Protamine messenger ribonucleic acid, testis, rainbow-trout, isolation and purification of (Gedamu, L., Iatrou, K. & Dixon, G. H.) 589–599
Protease, *see* Proteinase
Protein, glucocorticoid-receptor, cytosol, liver, rat, maximization of the purification of the complex of glucocorticoid and, by deoxyribonucleic acid–cellulose chromatography (Eisen, H. J. & Glinsmann, W. H.) 177–183
Protein, C-, interaction of, with heavy meromyosin and subfragment 2 of skeletal-muscle myosin (Starr, R. & Offer G.) 813–816
Protein kinase, adenosine 3′:5′-cyclic monophosphate-dependent, nucleoplasmic, liver, rat, histone-specific, partial purification and properties of (Neumann, J. R., O'Meara, A. R. & Herrmann, R. L.) 123–135
Protein, oestrogen-receptor, cytosol, mammary-gland, mouse, lactating, effect of heparin on the aggregation of, and interaction of the receptor protein with heparin–Sepharose (Auricchio, F., Rotondi, A., Sampaolo, P. & Schiavone, E.) 137–141
Protein–polysaccharides, *see* Proteoglycans
Protein, retinoic acid-binding, skin, chick-embryo, determination of binding affinities of retinoids to ox serum albumin and to (Sani, B. P., Titus, B. C. & Banerjee, C. K.) 711–717
Proteinase, alkaline, skeletal-muscle, rat, purification and properties of, associated with myofibrillar proteins (Dahlmann, B. & Reinauer, H.) 803–810
Proteinases, metallo-, neutral, bone, rabbit, two, separation of latent forms of collagenase and of, that when activated degrade gelatin and cartilage proteoglycans (Sellers, A., Reynolds, J. J. & Meikle, M. C.) 493–496
Protein–polysaccharides, *see* Proteoglycans
Proteins, criteria for recognition of the structure and micropolarity of pyridoxal 5′-phosphate-binding sites in (Cortijo, M., Jimenez, J. S. & Llor, J.) 497–500
Proteins, membrane, erythrocyte, human, cross-linking by periodate of, and distribution of intramembrane particles (Gahmberg, C. G., Virtanen, I. & Wartiovaara, J.) 683–686
Proteins, myofibrillar, skeletal-muscle, rat, purification and properties of an alkaline proteinase associated with (Dahlmann, B. & Reinauer, H.) 803–810
Proteins, simple sensitive method for the determination of polypeptides and, in polyacrylamide gels after isoelectric focusing (Storring, P. L. & Tiplady, R. J.) 79–82
Proteins, troponin-like, adductor-muscle, scallop, preparation and properties of (Goldberg, A. & Lehman, W.) 413–418
Proteoglycans, cartilage, laryngeal, pig, self-association of subunits of (Sheehan, J. K., Nieduszynski, I. A., Phelps, C. F., Muir, H. & Hardingham, T. E.) 109–114
Proteoglycans, cartilage, separation from rabbit bone of latent forms of collagenase and of two neutral metalloproteinases that when activated degrade gelatin and (Sellers, A., Reynolds, J. J. & Meikle, M. C.) 493–496
Proteoglycans, effect of vitamin D deficiency on, of chick growth cartilage (Dickson, I. R. & Roughley, P. J.) 675–682
Protons, exchange of, involving C-1 *pro-S* hydrogen atoms of dihydroxyacetone phosphate and fructose 1,6-bisphosphate catalysed by class-I and class-II fructose bisphosphate aldolases (Galdes, A. & Hill, H. A. O.) 539–542
Pseudomonas testosteroni N.C.I.B. 10808, purification and properties of urocanase from (Hacking, A. J., Bell, M. V. & Hassall, H.) 41–50
Pseudouridylate residues, evidence for lack of exact numerical correspondence between the contents of 2′-*O*-methyl groups and, of ribosomal ribonucleic acids from cells of three vertebrate species (Hughes, D. G. & Maden, B. E. H.) 781–786
Purple membrane, *see* Membrane, purple
Pyridoxal 5′-phosphate, criteria for recognition of the structure and micropolarity of binding sites for, in proteins (Cortijo, M., Jimenez, J. S. & Llor, J.) 497–500
Pyridoxal 5′-phosphate, reaction of amino-oxyacetate with enzymes dependent on (John, R. A., Charteris, A. & Fowler, L. J.) 771–779
Pyrophosphate, inorganic, effects of substrate, metal ions and template on the inhibition by inorganic phosphate and, of the activity of bacteriophage-Qβ ribonucleic acid polymerase (Brooks, R. R. & Andersen, J. A.) 725–732

1-Pyrrolidinecarbothioic acid (2-pyridylmethylene)hydrazide, simplified synthesis of, and its action as an inhibitor of phosphatases (Coburn, S. P. & Schaltenbrand, W. E.) 485–488
Pyruvate decarboxylase, intracellular location of citrate synthase, 3-hydroxyacyl-coenzyme A dehydrogenase and, in lactating rat mammary gland (Taylor, D. J., Crabtree, R. & Smith, G. H.) 273–275
Pyruvate, pathway for the metabolism of, to ethanol in *Entamoeba histolytica* (Lo, H.-S. & Reeves, R. E.) 225–230

Radish (*Raphanus sativus*) seedlings, effect of 2-(4-methyl-2,6-dinitroanilino)-*N*-methylpropionamide on the biosynthesis of ribosomal ribonucleic acid in (Aspart, L., Cooke, R., Delseny, M. & Guitton, Y.) 607–611
Rana pipiens, *see* Frog
Rana temporaria, *see* Frog
Raphanus sativus, *see* Radish
Red blood cells, *see* Erythrocytes
Reticulocytes, rabbit, characterization of monoferric fragments obtained by tryptic cleavage of ox transferrin and their inability to donate iron to (Brock, J. H., Arzabe, F. R., Richardson, N. E. & Deverson, E. V.) 73–78
Retinoic acid-binding protein, skin, chick-embryo, determination of binding affinities of retinoids to ox serum albumin and to (Sani, B. P., Titus, B. C. & Banerjee, C. K.) 711–717
Retinoids, determination of binding affinities of, to chick-embryo skin retinoic acid-binding protein and to ox serum albumin (Sani, B. P., Titus, B. C. & Banerjee, C. K.) 711–717
Rhodopseudomonas capsulata, location of *O*-methyl sugars in antigenic lipopolysaccharides and polysaccharides of various photosynthetic bacteria and cyanobacteria including (Tharanathan, R. N., Mayer, H. & Weckesser, J.) 403–408
Rhodopseudomonas gelatinosa, location of *O*-methyl sugars in antigenic lipopolysaccharides and polysaccharides of various photosynthetic bacteria and cyanobacteria including (Tharanathan, R. N., Mayer, H. & Weckesser, J.) 403–408
Ribonucleic acid, messenger, protamine, testis, rainbow-trout, isolation and purification of (Gedamu, L., Iatrou, K. & Dixon, G. H.) 589–599
Ribonucleic acid polymerase, bacteriophage-Qβ, effects of substrate, metal ions and template on the inhibition by inorganic phosphate and inorganic pyrophosphate of the activity of (Brooks, R. R. & Andersen, J. A.) 725–732
Ribonucleic acid, ribosomal, effect of 2-(4-methyl-2,6-dinitroanilino)-*N*-methylpropionamide on the biosynthesis of, in radish seedlings (Aspart, L., Cooke, R., Delseny, M. & Guitton, Y.) 607–611
Ribonucleic acid, transfer, *Escherichia coli*, use of chemical modification as a probe of conformational changes during aminoacylation of (Lowdon, M. & Goddard, J. P.) 601–606
Ribonucleic acids, ribosomal, evidence for lack of exact numerical correspondence between the contents of pseudouridylate residues and 2′-*O*-methyl groups of, from cells of three vertebrate species (Hughes, D. G. & Maden, B. E. H.) 781–786
Ribonucleic acids, ribosomal, nucleolar and nucleoplasmic, liver, rat, quantitatiye analysis of (Dabeva, M. D., Dudov, K. P., Hadjiolov, A. A. & Stoykova, A. S.) 367–374
Ribonucleic acids, ribosomal, processing and migration of, in the nucleoli and nucleoplasm of rat liver nuclei (Dudov, K. P., Dabeva, M. D., Hadjiolov, A. A. & Todorov, B. N.) 375–383
Ribosomal ribonucleic acid, *see* Ribonucleic acid, ribosomal
Ribulose bisphosphate carboxylase, comparison of the kinetic properties of, in extracts of chloroplasts from six higher-plant species utilizing the reductive pentose phosphate pathway (Delaney, M. E. & Walker, D. A.) 477–482
Rickets, effect of, on the proteoglycans of chick growth cartilage (Dickson, I. R. & Roughley, P. J.) 675–682

Saccharomyces cerevisiae, estimation of rate and dissociation constants involving ternary complexes in reactions catalysed by alcohol dehydrogenase from (Dickinson, F. M. & Dickenson, C. J.) 629–637
Saccharomyces cerevisiae, inhibition by ethanol, acetaldehyde and trifluoroethanol of reactions catalysed by alcohol dehydrogenases from horse liver and (Dickenson, C. J. & Dickinson, F. M.) 613–627
Salmo gairdnerii, *see* Trout, rainbow
Scallop (*Aequipecten irradians*), preparation and properties of troponin-like proteins from adductor muscle of (Goldberg, A. & Lehman, W.) 413–418
Seedlings, radish, effect of 2-(4-methyl-2,6-dinitroanilino)-*N*-methylpropionamide on the biosynthesis of ribosomal ribonucleic acid in (Aspart, L., Cooke, R., Delsey, M. & Guitton, Y.) 607–611
Seeds, lentil and pea, isolation and characterization of phytoferritins from (Crichton, R. R., Ponce-Ortiz, Y., Koch, M. H. J., Parfait, R. & Stuhrmann, H. B.) 349–356
Seminal plasma, human, purification and properties of two plasminogen activators from (Propping, D., Zaneveld, L. J. D., Tauber, P. F. & Schumacher, G. F. B.) 435–444
Serum albumin, *see* Albumin, serum
Skeletal muscle, *see* Muscle, skeletal
Skin, chick-embryo, determination of binding affinities of retinoids to ox serum albumin and to retinoic acid-binding protein from (Sani, B. P., Titus, B. C. & Banerjee, C. K.) 711–717
Slime mould (*Physarum polycephalum*), heterogeneity of cytoplasmic deoxyribonucleic acid polymerase from (Zänker, K. S. & Schiebel, W.) 445–451
Solanum tuberosum, *see* Potato
Spectrin, membrane, erythrocyte, human, cross-linking by periodate of, and other membrane proteins and distribution of intramembrane particles (Gahmberg, C. G., Virtanen, I. & Wartiovaara, J.) 683–686
Spermatozoa, salmon, relative reactivity of the phosphodiester site thymidylyl(3′-5′)thymidine of deoxyribonucleic acid from, towards alkylation by dimethyl sulphate, ethyl methanesulphonate, *N*-ethyl-*N*-nitrosourea and *N*-methyl-*N*-nitrosourea (Swenson, D. H. & Lawley, P. D.) 575–587

Spinach (*Spinacia oleracea*) leaves, comparison of the kinetic properties of ribulose bisphosphate carboxylase in extracts of chloroplasts from, and from five other higher-plant species utilizing the reductive pentose phosphate pathway (Delaney, M. E. & Walker, D. A.) 477–482
Spinacia oleracea, *see* Spinach
Spleen, horse, incorporation and release of iron and inorganic phosphate in ferritin from (Treffry, A. & Harrison, P. M.) 313–320
Substance P, inactivation of, in rat kidney (Ward, P. E. & Johnson, A. R.) 143–148
Sunflower (*Helianthus annuus*) leaves, comparison of the kinetic properties of ribulose biphosphate carboxylase in extracts of chloroplasts from, and from five other higher-plant species utilizing the reductive pentose phosphate pathway (Delaney, M. E. & Walker, D. A.) 477–482
Supernatant fraction, *see* Cytosol

Testis, rainbow-trout, isolation and purification of protamine messenger ribonucleic acid from (Gedamu, L., Iatrou, K. & Dixon, G. H.) 589–599
Tetrahydrobiopterin, kinetics of the biosynthesis of, by rabbit brain tetrahydrofolate dehydrogenase (Abelson, H. T., Spector, R., Gorka, C. & Fosburg, M.) 267–268
Tetrahydrofolate dehydrogenase, brain, rabbit, kinetics of the biosynthesis of tetrahydrobiopterin by (Abelson, H. T., Spector, R., Gorka, C. & Fosburg, M.) 267–268
Tetrahydrofolate dehydrogenase, *Lactobacillus casei*, use of ultraviolet difference spectroscopy for study of the binding of substrate and inhibitors to (Hood, K. & Roberts, G. C. K.) 357–366
3α,7α,12α,22ξ-Tetrahydroxy-5β-cholestan-26-oic acid, identification of the taurine conjugate of, as a major component of green-turtle bile salts (Haslewood, G. A. D., Ikawa, S., Tökés, L. & Wong, D.) 409–412
Thiol group, active-site, use of two-protonic-state electrophiles as reactivity probes for, of papain (Shipton, M. & Brocklehurst, K.) 385–401
Thiol group, reactive, use of a chromophoric reporter group to study the role of, of rabbit skeletal-muscle creatine kinase (Keighren, M. A. & Price, N. C.) 269–272
Thymidylyl(3′-5′)thymidine, phosphodiester-site, relative reactivity of, of salmon sperm deoxyribonucleic acid towards alkylation by dimethyl sulphate, ethyl methanesulphonate, *N*-ethyl-*N*-nitrosourea and *N*-methyl-*N*-nitrosourea (Swenson, D. H. & Lawley, P. D.) 575–587
Transfer ribonucleic acid, *see* Ribonucleic acid, transfer
Transferrin, ox, characterization of monoferric fragments obtained by tryptic cleavage of, and their inability to donate iron to rabbit reticulocytes (Brock, J. H., Arzabe, F. R., Richardson, N. E. & Deverson, E. V.) 73–78
Transplantation antigens, human (HLA), incorporation of, into liposomes (Turner, M. J. & Sanderson, A. R.) 505–508
Trichoderma koningii I.M.I. 73022, purification and properties of four endoglucanase components of the cellulase complex of, with special reference to their action on cellulose when acting alone and in synergism with the cellobiohydrolase component (Wood, T. M. & McCrae, S. I.) 61–72
Trifluoroethanol, inhibition by ethanol, acetaldehyde and, of reactions catalysed by yeast and horse liver alcohol dehydrogenases (Dickenson, C. J. & Dickinson, F. M.) 613–627
3α,12α,22ξ-Trihydroxy-5β-cholestan-26-oic acid, identification of the taurine conjugate of, as a major component of green-turtle bile salts (Haslewood, G. A. D., Ikawa, S., Tökés, L. & Wong, D.) 409–412
Troponin I, skeletal-muscle and heart-muscle, human, properties of, and from the muscles of other mammalian species (Cummins, P. & Perry, S. V.) 251–259
Troponins, adductor-muscle, scallop, preparation and properties of (Goldberg, A. & Lehman, W.) 413–418
Trout, rainbow (*Salmo gairdnerii*), isolation and purification of protamine messenger ribonucleic acid from testis of (Gedamu, L., Iatrou, K. & Dixon, G. H.) 589–599
Tubers, potato, properties of the lectin from, and the nature of its glycoprotein linkages (Allen, A. K., Desai, N. N., Neuberger, A. & Creeth, J. M.) 665–674
Turkey, stopped-flow spectrophotometric studies on the reaction of xanthine dehydrogenase from the liver of, with reducing substrates (Ní Fhaoláin, Í., Hynes, M. J. & Coughlan, M. P.) 83–88

Uric acid, electron-paramagnetic-resonance spectroscopy of complexes of milk xanthine oxidase with xanthine and (Bray, R. C., Barber, M. J., & Lowe D. J.) 653–658
Uridine diphosphate glucuronosyltransferases, microsomal, liver, rabbit, separation of the activities of, specific for oestrone and *p*-nitrophenol (Tukey, R. H., Billings, R. E. & Tephly, T. R.) 659–663
Uridine diphosphate glucuronyltransferase, microsomal, liver, rat, purified, content of phospholipids and activity of (Burchell, B. & Hallinan, T.) 821–824
Urine, human, simple high-yield procedure for isolation of kallikreins from (Oza, N. B. & Ryan, J. W.) 285–288
Urocanase, *Pseudomonas testosteroni* N.C.I.B. 10808, purification and properties of (Hacking, A. J., Bell, M. V. & Hassall, H.) 41–50
Urokinase, evidence for non-specific dissociable binding obtained during investigation by isoelectric focusing of the interaction of human α_2-macroglobulin with (Vahtera, E. & Hamberg, U. 767–770

Vegetable-marrow (*Cucurbita pepo*) leaves, electron-paramagnetic-resonance signals from the iron–sulphur centre and haem group during turnover of nitrite reductase from (Cammack, R., Hucklesby, D. P. & Hewitt, E. J.) 519–526
Vitamin D, effect of deficiency of, on the proteoglycans of chick growth cartilage (Dickson, I. R. & Roughley, P. J.) 675–682

Xanthine dehydrogenase, liver, turkey, stopped-flow spectrophotometric studies on the reaction of, with reducing substrates (Ní Fhaoláin, Í., Hynes, M. J. & Coughlan, M. P.) 83–88
Xanthine, electron-paramagnetic-resonance spectroscopy of complexes of milk xanthine oxidase with uric acid and (Bray, R. C., Barber, M. J. & Lowe, D. J.) 653–658

Xanthine oxidase, milk, electron-paramagnetic-resonance spectroscopy of complexes of, with xanthine and uric acid (Bray, R. C., Barber, M. J. & Lowe, D. J.) 653–658

Yeast (*Saccharomyces cerevisiae*), estimation of rate and dissociation constants involving ternary complexes in reactions catalysed by alcohol dehydrogenase from (Dickinson, F. M. & Dickenson, C. J.) 629–637

Yeast (*Saccharomyces cerevisiae*), inhibition by ethanol, acetaldehyde and trifluoroethanol of reactions catalysed by alcohol dehydrogenases from horse liver and Dickenson, C. J. & Dickinson, F. M.) 613–627

Zinc ions, simplified synthesis of 1-pyrrolidinecarbothioic acid (2-pyridylmethylene)hydrazide and its action as a selective chelator of, and inhibitor of phosphatases (Coburn, S. P. & Schaltenbrand, W. E.) 485–488

JECTS

ethanol,
atalysed
er and
7

pothioic
ion as a
phatases
488